

BROWDY AND NEIMARK, PLLC.

ATTORNEYS AT LAW  
PATENT AND TRADEMARK CAUSES

SUITE 300

419 SEVENTH STREET, N W  
WASHINGTON, D. C 20004-2299

TELEPHONE (202)-628-5197

SHERIDAN NEIMARK  
ROGER L. BROWDY **52222 U.S. PTO**

ANNE M. KORNBAUM  
NORMAN J. LATKIN  
NICK BROMER\*  
(\*PA BAR ONLY)

OF COUNSEL  
IVER P. COOPER

TELECOPIER FACSIMILE  
(202) 737-3528  
(202) 393-1012

E-MAIL  
BrwdyNmrmk@nmaa.org

SENIOR COUNSEL  
ALVIN BROWDY

PATENT AGENT  
ALLEN C. YUN, PH.D.

February 20, 1997

Hon. Commissioner of Patents  
and Trademarks  
Washington, D.C. 20231

RE: New Patent Application in U.S.  
Robert CAMPBELL et al.  
Atty's Docket: CAMPBELL=2A

Sir:

Transmitted herewith is a patent application entitled HYBRID PROTEINS. The inventors are Robert CAMPBELL, Bradford A. JAMESON and Scott C. CHAPPEL. Priority under 35 USC §119 is claimed from U.S. Provisional Application no. 60/011,936 filed 20 February 1996.

Attached are the specification (49 pages), including 20 claims (1 independent) and an abstract, plus 7 sheets of drawings (Figures 1-6), and a return postcard. In accordance with 37 CFR 1.53(a) and (b), it is respectfully requested that a serial number and filing date be assigned to this application as of the date of receipt of the present papers. In accordance with the present procedures of the U.S. Patent and Trademark Office, an executed Declaration and the filing fee for the present application will be filed in due course.<sup>1</sup>

<sup>1</sup> No authorization is given for charging the filing fee at the present time. However, at such time that the declaration is filed, but not before, you are authorized to charge whatever excess fees are necessary (including the filing fee and any extension of time fees then due) to Deposit Account 02-4035, if any such fees due are not fully covered by check filed at that time.

In re of CAMPBELL=2A

The attached specification includes a paper copy Sequence Listing section according to 37 CFR §1.821(c) as pages 32-45. Also attached hereto is a 3 1/2" floppy disk containing the "Sequence Listing" in computer readable form in accordance with 37 CFR §1.821(e). I hereby state, in accordance with 37 CFR §1.821(f), that the content of the paper and computer readable copies of the sequence listing, submitted in accordance with 37 CFR §1.821(c) and (e), respectively, are believed to be the same.


The attorneys of record in this application will be Sheridan Neimark, Reg. No. 20,520; Roger L. Browdy, Reg. No. 25,618; Anne M. Kornbau, Reg. No. 25,884; Norman J. Latker, Reg. No. 19,963; Iver P. Cooper, Reg. No. 28,005; \*Allen C. Yun, Reg. No. 37,971 and Nick Bromer, Reg. No. 33,478 (\*Patent Agent). Please send all correspondence with respect to this case to:

BROWDY AND NEIMARK, P.L.L.C.  
419 Seventh Street, N.W.  
Washington, D.C. 20004

Please direct all telephone calls to Browdy and Neimark at (202) 628-5197.

Respectfully submitted,  
BROWDY AND NEIMARK, P.L.L.C.

By

  
Roger L. Browdy  
Registration No. 25,618

RLB:bcs

f:\user3\97feb\cam2a.53

2/

62222 U.S. PTO



HYBRID PROTEINS

A/NO  
File

084804166

02/20/97  
CROSS REFERENCE TO RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application no. 60/011,936, filed February 20, 1996.

FIELD OF THE INVENTION

The present invention relates to a hybrid protein comprising two coexpressed amino acid sequences forming a dimer, each comprising:

a) at least one amino acid sequence selected from a homomeric receptor, a chain of a heteromeric receptor, a ligand, and fragments thereof; and

b) a subunit of a heterodimeric proteinaceous hormone or fragments thereof; in which (a) and (b) are bonded directly or through a peptide linker, and, in each couple, the two subunits (b) are different and capable of aggregating to form a dimer complex.

BACKGROUND OF THE INVENTION

Protein-protein interactions are essential to the normal physiological functions of cells and multicellular organisms. Many proteins in nature exhibit novel or optimal functions when complexed with one or more other protein chains. This is illustrated by various ligand-receptor combinations that contribute to regulation of cellular activity. Certain ligands, such as tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), TNF $\beta$ , or human chorionic gonadotropin (hCG), occur as multi-subunit complexes. Some of these complexes contain multiple copies of the same subunit. TNF $\alpha$  and TNF $\beta$  (collectively referred to hereafter as TNF) are homotrimers formed by three identical subunits (1-4). Other ligands are composed of non-identical subunits. For example, hCG is a heterodimer (5-7). Receptors may also occur or function as multi-chain complexes. For example, receptors for TNF transduce a signal after being aggregated to form dimers (8,9). Ligands to these receptors

promote aggregation of two or three receptor chains, thereby affording a mechanism of receptor activation. For example, TNF-mediated aggregation activates TNF receptors (10-12).

The modulation of protein-protein interactions can be a useful mechanism for therapeutic intervention in various diseases and pathologies. Soluble binding proteins, that can interact with ligands, can potentially sequester the ligand away from the receptor, thereby reducing the activation of that particular receptor pathway. Alternatively, sequestration of the ligand may delay its elimination or degradation, thereby increasing its duration of effect, and perhaps its apparent activity *in vivo*. In the case of TNF, soluble TNF receptors have been primarily associated with inhibition of TNF activity (13-17).

Soluble binding proteins may be useful for treating human diseases. For example, soluble TNF receptors have been shown to have efficacy in animal models of arthritis (18,19).

Since TNF has three binding sites for its receptor (10-12), and dimerization of the cell surface receptor is sufficient for bioactivity (8,9), it is likely that binding of a single soluble receptor to TNF will leave open the possibility that this 1:3 complex of soluble receptor:TNF (trimer) can still bind and activate a pair of cell surface TNF receptors. To achieve an inhibitory effect, it would be expected that two of the receptor binding sites on the TNF trimer must be occupied or blocked by the soluble binding protein. Alternatively, the binding protein could block proper orientation of TNF at the cell surface.

Generally speaking, the need was felt of synthesizing proteins that contain two receptor (or ligands) chains, as dimeric hybrid protein. See Wallach et al., U.S. patent 5,478,925.

The primary strategy employed for generating dimeric or multimeric hybrid proteins, containing binding domains from extracellular receptors, has been to fuse these proteins to the constant regions of an antibody heavy chain.

This strategy led, for example, to the construction of CD4 immunoadhesins (20). These are hybrid molecules consisting of the first two (or all four) immunoglobulin-like domains of CD4 fused to the constant region of antibody heavy and light chains. This strategy for creating hybrid molecules was adapted to the receptors for TNF (10,16,21) and led to the generation of constructs with higher *in vitro* activity than the monomeric soluble binding proteins.

It is widely held that the higher *in vitro* potency of the dimeric fusion proteins should translate into higher *in vivo* activity. One study does support this, revealing an at least 50-fold higher activity for a p75(TBP2)-Ig fusion protein in protecting mice from the consequences of intravenous LPS injection (16).

However, despite the widespread utilization of immunoglobulin fusion proteins, this strategy has several drawbacks. One is that certain immunoglobulin Fc domains participate in effector functions of the immune system. These functions may be undesirable in a particular therapeutic setting (22).

A second limitation pertains to the special cases where it is desirable to produce heteromeric fusion proteins, for example soluble analogs of the heteromeric IL-6 or type I interferon receptors. Although there are numerous methods for producing bifunctional antibodies (e.g., by co-transfection or hybridoma fusions), the efficiency of synthesis is greatly compromised by the mixture of homodimers and heterodimers that typically results (23). Recently there have been several reports describing the use of leucine zipper motifs to guide assembly of heterodimers (24-26). This appears to be a promising approach for research purposes, but the non-native or intracellular sequences employed may not be suitable for chronic applications in the clinic due to antigenicity. The efficiency of assembly and stability post assembly may also be limitations.

On the other hand, in the particular case of TNF receptors, certain modifications to the p55 TNF receptor have

been found to facilitate homodimerization and signaling in the absence of ligand (27,28). It has been found that a cytoplasmic region of the receptor, termed the "death domain," can act as a homodimerization motif (28,30). As an alternative to an immunoglobulin hybrid protein, fusion of the extracellular domain of the TNF receptor to its cytoplasmic death domain could conceivably result in a secreted protein which can dimerize in the absence of TNF. Such fusion proteins have been disclosed and claimed in the International Patent Application WO 95/31544.

A third further strategy employed for generating dimers of soluble TNF receptors has been to chemically cross-link the monomeric proteins with polyethylene glycol (31).

#### SUMMARY OF THE INVENTION

An alternative for obtaining such dimeric proteins, offering some important advantages, is the one of the present invention and consists in using a natural heterodimeric scaffold corresponding to a circulating non-immunoglobulin protein with a long half-life. A preferred example is hCG, a protein that is secreted well, has good stability, and has a long half-life (32-33). Given hCG's prominent role as a marker of pregnancy, many reagents have been developed to quantitate and study the protein *in vitro* and *in vivo*. In addition, hCG has been extensively studied using mutagenesis, and it is known that small deletions to the protein, such as removal of five residues at the extreme carboxyl-terminus of the  $\alpha$  subunit, can effectively eliminate its biological activity while preserving its capability to form heterodimer (34,35). Small insertions, of up to 30 amino acids, have been shown to be tolerated at the amino- and carboxyl-termini of the  $\alpha$  subunit (36), while fusion of the  $\alpha$  subunit to the carboxyl terminus of the  $\beta$  subunit also had little effect on heterodimer formation (37).

An analog of hCG in which an immunoglobulin Fc domain was fused to the C-terminus of hCG  $\beta$  subunit has also been

reported; however, this construct was not secreted and no effort was made to combine it with an  $\alpha$  subunit (38).

Therefore, the main object of the present invention is a hybrid protein comprising two coexpressed amino acid sequences forming a dimer, each comprising:

a) at least one amino acid sequence selected among a homomeric receptor, a chain of a heteromeric receptor, a ligand, and fragments thereof; and

b) a subunit of a heterodimeric proteinaceous hormone, or fragments thereof; in which (a) and (b) are bonded directly or through a peptide linker, and in each couple the two subunits (b) are different and capable of aggregating forming a dimer complex.

According to the present invention, the linker may be enzymatically cleavable.

Sequence (a) is preferably selected among: the extracellular domain of the TNF Receptor 1 (55 kDa, also called TBP1), the extracellular domain of the TNF Receptor 2 (75 kDa, also called TBP2), or fragments thereof still containing the ligand binding domain; the extracellular domains of the IL-6 receptors (also called gp80 and gp130); the extracellular domain of the IFN  $\alpha/\beta$  receptor or IFN  $\gamma$  receptor; a gonadotropin receptor or its extracellular fragments; antibody light chains, or fragments thereof, optionally associated with the respective heavy chains; antibody heavy chains, or fragments thereof, optionally associated with the respective light chains; antibody Fab domains; or ligand proteins, such as cytokines, growth factors or hormones other than gonadotropins, specific examples of which include IL-6, IFN- $\beta$ , TPO, or fragments thereof.

Sequence (b) is preferably selected among a hCG, FSH, LH, TSH, inhibin subunit, or fragments thereof.

Modifications to the proteins, such as chemical or protease cleavage of the protein backbone, or chemical or enzymatic modification of certain amino acid side chains, can be used to render the components of the hybrid protein of the invention inactive. This restriction of activity may also be

accomplished through the use of recombinant DNA techniques to alter the coding sequence for the hybrid protein in a way that results directly in the restriction of activity to one component, or that renders the protein more amenable to subsequent chemical or enzymatic modification.

The above hybrid proteins will result in monofunctional, bifunctional or multifunctional molecules, depending on the amino acid sequences (a) that are combined with (b). In each couple, (a) can be linked to the amino termini or to the carboxy termini of (b), or to both.

A monoclonal hybrid protein of the present invention can, for instance, comprise the extracellular domain of a gonadotropin receptor linked to one of the corresponding receptor-binding gonadotropin subunits. According to such an embodiment, the hybrid protein of the invention can be a molecule in which, for example, the FSH receptor extracellular domain is linked to FSH to increase plasma half-life and improve biological activity.

This preparation can be employed to induce follicular maturation in assisted reproduction methods, such as ovulation induction or *in vitro* fertilisation, and to serve as a means to dramatically amplify the biological activity of the hormone essential for the success of the process, thus reducing the requirement for both the hormone itself and the number of injections to achieve ovulation.

The FSH receptor and the production of the extracellular domain of the human FSH receptor have been described respectively in WO 92/16620 and WO 96/38575.

According to a particular embodiment, the extracellular domain of the FSH receptor (ECD) can be fused in frame with a peptide linker that contains the thrombin recognition/cleavage site (29) and represents a "tethered" arm. The peptide linker links the extracellular domain of FSH with a FSH subunit. This will allow for removal of the extracellular domain of the FSH receptor by cleavage at the thrombin cleavage site as the molecule comes in contact with thrombin in the systemic circulation.



In another embodiment, instead of the thrombin cleavage site, an enzyme recognition site for an enzyme that is found in greatest abundance in the ovary is used. In this way, as the ECD-FSH molecule travels to the ovary, it will be exposed to enzymes found in the highest concentrations in that tissue and the ECD will be removed so that the FSH can interact with the membrane bound receptor.

In yet another embodiment, instead of an enzyme recognition site, a flexible hinge region is cloned between ECD and FSH so that the ECD will not be enzymatically removed from the hormone. In this way, when the ECD-FSH molecule arrives at the ovary, a competition will be established between the hinge-attached ECD and the ECD of the FSH receptor found on the ovarian cell membrane.

In a further preferred embodiment of the invention, the hybrid protein consists of the aggregation between a couple of aa sequences, one of which contains TBP1 (or the fragments from aa 20 to aa 161 or to aa 190) as (a) and the  $\alpha$  subunit of hCG as (b), and the other contains always TBP1 (or the same fragments as above) as (a) and the  $\beta$  subunit of hCG, or fragments thereof, as (b). According to this embodiment, depending on the particular sequence that is chosen as (b) (the entire  $\beta$  subunit of hCG, or fragments or modifications thereof), the resulting hybrid protein will have one activity (only that of TBP1) or a combination of activities (that of TBP1 with that of hCG). In this latter case the hybrid protein can be used, for example, in the combined treatment of Kaposi's sarcoma and metabolic wasting in AIDS.

In a further embodiment of the invention, one or more covalent bonds between the two subunits (b) are added to enhance the stability of the resulting hybrid protein. This can be done, e.g., by adding one or more non-native interchain disulfide bonds. The sites for these cross-links can be deduced from the known structures of the heterodimeric hormones. For example, a suitable site in hCG could be to place cysteine residues at  $\alpha$  subunit residue Lys45 and  $\beta$  subunit residue Glu21, replacing a salt bridge (non-covalent

bond) with a disulfide bond (covalent bond). Another object of the present invention are PEGylated or other chemically modified forms of the hybrid proteins.

A further object of the present invention is a DNA molecule comprising the DNA sequence coding for the above hybrid protein, as well as nucleotide sequences substantially the same. "Nucleotide sequences substantially the same" includes all other nucleic acid sequences which, by virtue of the degeneracy of the genetic code, also code for the given amino acid sequence.

For the production of the hybrid protein of the invention, the DNA sequence (a) is obtained from existing clones, as is (b). The DNA sequence coding for the desired sequence (a) is ligated with the DNA sequence coding for the desired sequence (b). Two of these fused products are inserted and ligated into a suitable plasmid or each into a different plasmid. Once formed, the expression vector, or the two expression vectors, is introduced into a suitable host cell, which then expresses the vector(s) to yield the hybrid protein of the invention as defined above.

The preferred method for preparing the hybrid of the invention is by way of PCR technology using oligonucleotides specific for the desired sequences to be copied from the clones encoding sequences (a) and (b).

Expression of any of the recombinant proteins of the invention as mentioned herein can be effected in eukaryotic cells (e.g., yeasts, insect or mammalian cells) or prokaryotic cells, using the appropriate expression vectors. Any method known in the art can be employed.

For example the DNA molecules coding for the proteins obtained by any of the above methods are inserted into appropriately constructed expression vectors by techniques well known in the art (see Sambrook et al, 1989). Double stranded cDNA is linked to plasmid vectors by homopolymeric tailing or by restriction linking involving the use of synthetic DNA linkers or blunt-ended ligation techniques: DNA ligases are

used to ligate the DNA molecules and undesirable joining is avoided by treatment with alkaline phosphatase.

In order to be capable of expressing the desired protein, an expression vector should comprise also specific nucleotide sequences containing transcriptional and translational regulatory information linked to the DNA coding the desired protein in such a way as to permit gene expression and production of the protein. First in order for the gene to be transcribed, it must be preceded by a promoter recognizable by RNA polymerase, to which the polymerase binds and thus initiates the transcription process. There are a variety of such promoters in use, which work with different efficiencies (strong and weak promoters).

For eukaryotic hosts, different transcriptional and translational regulatory sequences may be employed, depending on the nature of the host. They may be derived from viral sources, such as adenovirus, bovine papilloma virus, Simian virus or the like, where the regulatory signals are associated with a particular gene which has a high level of expression. Examples are the TK promoter of the Herpes virus, the SV40 early promoter, the yeast gal4 gene promoter, etc. Transcriptional initiation regulatory signals may be selected which allow for repression and activation, so that expression of the genes can be modulated.

The DNA molecule comprising the nucleotide sequence coding for the hybrid protein of the invention is inserted into a vector(s), having the operably linked transcriptional and translational regulatory signals, which is capable of integrating the desired gene sequences into the host cell. The cells which have been stably transformed by the introduced DNA can be selected by also introducing one or more markers which allow for selection of host cells which contain the expression vector. The marker may also provide for phototrophy to a auxotrophic host, biocide resistance, e.g., antibiotics, or heavy metals such as copper, or the like. The selectable marker gene can either be directly linked to the DNA gene sequences to be expressed, or introduced into the same cell by

co-transfection. Additional elements may also be needed for optimal synthesis of proteins of the invention.

Factors of importance in selecting a particular plasmid or viral vector include: the ease with which recipient cells that contain the vector may be recognized and selected from those recipient cells which do not contain the vector; the number of copies of the vector which are desired in a particular host; and whether it is desirable to be able to "shuttle" the vector between host cells of different species.

Once the vector(s) or DNA sequence containing the construct(s) has been prepared for expression, the DNA construct(s) may be introduced into an appropriate host cell by any of a variety of suitable means: transformation, transfection, conjugation, protoplast fusion, electroporation, calcium phosphate-precipitation, direct microinjection, etc.

Host cells may be either prokaryotic or eukaryotic. Preferred are eukaryotic hosts, e.g., mammalian cells, such as human, monkey, mouse, and Chinese hamster ovary (CHO) cells, because they provide post-translational modifications to protein molecules, including correct folding or glycosylation at correct sites. Also, yeast cells can carry out post-translational peptide modifications including glycosylation. A number of recombinant DNA strategies exist which utilize strong promoter sequences and high copy number of plasmids which can be utilized for production of the desired proteins in yeast. Yeast recognizes leader sequences on cloned mammalian gene products and secretes peptides bearing leader sequences (i.e., pre-peptides).

After the introduction of the vector(s), the host cells are grown in a selective medium, which selects for the growth of vector-containing cells. Expression of the cloned gene sequence(s) results in the production of the desired proteins.

Purification of the recombinant proteins is carried out by any one of the methods known for this purpose, i.e., any conventional procedure involving extraction, precipitation, chromatography, electrophoresis, or the like. A further

purification procedure that may be used in preference for purifying the protein of the invention is affinity chromatography using monoclonal antibodies which bind the target protein and which are produced and immobilized on a gel matrix contained within a column. Impure preparations containing the recombinant protein are passed through the column. The protein will be bound to the column by the specific antibody while the impurities will pass through. After washing, the protein is eluted from the gel by a change in pH or ionic strength.

The term "hybrid protein", as used herein, generically refers to a protein which contains two or more different proteins or fragments thereof.

As used herein, "fusion protein" refers to a hybrid protein, which consists of two or more proteins, or fragments thereof, linked together covalently.

The term "aggregation", as used herein, means the formation of strong specific non-covalent interactions between two polypeptide chains forming a complex, such as those existing between the  $\alpha$  and  $\beta$  subunit of a heterodimeric hormone (such as FSH, LH, hCG or TSH).

The terms "ligand" or "ligand protein", as used herein, refer to a molecule, other than an antibody or an immunoglobulin, capable of being bound by the ligand-binding domain of a receptor; such molecule may occur in nature, or may be chemically modified or chemically synthesised.

The term "ligand-binding domain", as used herein, refers to a portion of the receptor that is involved in binding a ligand and is generally a portion or essentially all of the extracellular domain.

The term "receptor", as used herein, refers to a membrane protein, whose binding with the respective ligand triggers secondary cellular responses that result in the activation or inhibition of intracellular process.

In a further aspect, the present invention provides the use of the hybrid protein as a medicament. The medicament is preferably presented in the form of a pharmaceutical

composition comprising the protein of the invention together with one or more pharmaceutically acceptable carriers and/or excipients. Such pharmaceutical compositions represent yet a further aspect of the invention.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

The invention will be better understood by reference to the appended drawings, in which:

Figures 1(a) and 1(b) show the TBP(20-161)-hCG $\alpha$  and TBP(20-161)-hCG $\beta$  constructs, respectively, and the corresponding sequences (SEQ ID NOS:1-4).

Figures 2(a) and 2(b) show the TBP(20-190)-hCG $\alpha$  and TBP(20-190)-hCG $\beta$  constructs, respectively, and the corresponding sequences (SEQ ID NOS:5-8).

Figure 3 is a schematic summary of the constructs of Figures 1 and 2 showing p55 TNFR1, TBP1 and TBP1 fusion constructs. The linker sequences shown on the last two lines are SEQ ID NO:9 (Ala-Gly-Ala-Ala-Pro-Gly) and SEQ ID NO:10 (Ala-Gly-Ala-Gly).

Figure 4 is a graph illustrating the dose dependent protective effect of CHO cell expressed TBP-hCG(20-190) on TNF $\alpha$ -induced cytotoxicity on BT-20 cells and various controls.

Figure 5 is a graph illustrating the dose dependent protective effect of COS cell expressed TBP-hCG(20-190) on TNF $\alpha$ -induced cytotoxicity on BT-20 cells and various controls.

Figure 6 is a graph illustrating the dose dependent protective effect of affinity purified CHO cell expressed TBP-hCG(20-161) on TNF $\alpha$ -induced cytotoxicity on BT-20 cells and various controls.

#### **DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS**

The invention will now be described by means of the following Examples, which should not be construed as in any way limiting the present invention.

## EXAMPLES

### **Materials and Methods**

Cell lines used in this study were obtained from the American Type Culture Collection (ATCC), Rockville, Maryland, unless otherwise specified. The CHO-DUKX cell line was obtained from L. Chasin at Columbia University through D. Houseman at MIT (39). The CHO-DUKX cells, which lack a functional gene for dihydrofolate reductase, were routinely maintained in complete  $\alpha$ -plus Modified Eagles Medium ( $\alpha$ (+)MEM) supplemented with 10% fetal bovine serum (FBS). The COS-7 cells were routinely maintained in Dulbecco's Modified Eagles Medium (DMEM) supplemented with 10% FBS. Unless specified otherwise, cells were split to maintain them in log phase of growth, and culture reagents were obtained from GIBCO (Grand Island, New York).

#### 1. Assembly of the genetic constructs encoding the hybrid proteins

The numbering assignments for the p55 TNF receptor are based on the cloning paper from Wallach (40), while the numbering assignments for the hCG subunits are based on the numbering assignments from the Fiddes cloning papers (41,42). The designation TBP, or TNF binding protein, refers to the extracellular domain portions of the TNF receptors capable of binding TNF. In these Examples, the DNA constructs will be named as TBP-hybrid proteins, with the partner and region of TBP indicated in the construct nomenclature. All of the TBP-hCG constructs contain the human growth hormone (hGH) signal peptide in place of the native p55 signal sequence. In addition, the hGH signal peptide has been placed so that it immediately precedes TBP residue Asp20, which is anticipated to make this the first residue in the mature, secreted protein. These modifications are not essential to the basic concept of using hCG as a partner of the hybrid protein.

The DNAs encoding the hybrid proteins were constructed using PCR methodology (43).

a. TBP1(20-161)-hCG

The initial TBP-hCG construct was engineered to contain the ligand binding domain from the extracellular region of the p55 TNF receptor (from Asp20 inclusive of residue Cys161) fused though a short linker to the hCG  $\alpha$  and  $\beta$  subunits (starting at residues  $\alpha$ Cys7 or  $\beta$ Pro7, respectively). This construct, hereafter referred to as TBP1(20-161)-hCG, is a heterodimer of two modified hCG subunits, TBP1(20-161)-hCG $\alpha$  and TBP1(20-161)-hCG $\beta$ .

The oligodeoxynucleotide primers used for the TBP1(20-161)-hCG $\alpha$  construct were:

primer 1( $\alpha\beta$ ) TTT TCT CGA GAT GGC TAC AGG TAA GCG  
CCC (SEQ ID NO:11)  
primer 2( $\alpha$ ) ACC TGG GGC AGC ACC GGC ACA GGA GAC ACA  
CTC GTT TTC (SEQ ID NO:12)  
primer 3( $\alpha$ ) TGT GCC GGT GCT GCC CCA GGT TGC CCA GAA  
TGC ACG CTA CAG (SEQ ID NO:13)  
primer 4( $\alpha$ ) TTT TGG ATC CTT AAG ATT TGT GAT AAT AAC  
AAG TAC (SEQ ID NO:14)

These and all of the other primers described in these Examples were synthesized on an Applied Biosystems Model 392 DNA synthesis machine (ABI, Foster City, California), using phosphoramidite chemistry.

Since both of the TBP-hCG subunit constructs have the same 5'-end (i.e., the 5'-end of the hGH/TBP construct), primer 1( $\alpha\beta$ ) was used for both TBP-hCG subunit constructs. The other primers used for the TBP1(20-161)-hCG $\beta$  construct were:

primer 2( $\beta$ ) CCG TGG ACC AGC ACC AGC ACA GGA GAC  
ACA CTC GTT TTC (SEQ ID NO:15)  
primer 3( $\beta$ ) TGT GCT GGT GCT GGT CCA CGG TGC CGC  
CCC ATC AAT (SEQ ID NO:16)  
primer 4( $\beta$ ) TTT TGG ATC CTT ATT GTG GGA GGA TCG  
GGG TG (SEQ ID NO:17)

Primers 2( $\alpha$ ) and 3( $\alpha$ ) are reverse complements, and cover both the 3'-end of the coding region for the p55 extracellular domain, and the 5'-end of the hCG  $\alpha$  subunit. Similarly, primers 2( $\beta$ ) and 3( $\beta$ ) are also reverse



complements, and cover both the 3'-end of the coding region for the p55 extracellular domain, and the 5'-end of the hCG  $\beta$  subunit.

Two PCR reactions were run for each of the two TBP-hCG subunit constructs. The first used primers 1( $\alpha\beta$ ) and 2( $\alpha$  or  $\beta$ ), and used as the template a plasmid encoding soluble p55 residues 20-180 preceded by the hGH signal peptide (plasmid pCMVhGHspcDNA.pA4). The second used primers 3( $\alpha$  or  $\beta$ ) and 4( $\alpha$  or  $\beta$ ), and used as the template either plasmid pSVL-hCG $\alpha$  or pSVL-hCG $\beta$  (44). The PCR was performed using Vent (TM) polymerase from New England Biolabs (Beverly, Massachusetts) in accordance with the manufacturer's recommendations, using for each reaction 25 cycles and the following conditions:

100  $\mu$ g of template DNA

1  $\mu$ g of each primer

2U of Vent(TM) polymerase (New England Biolabs)

denaturation at 99°C for 30 seconds

annealing at: 59°C for 30 seconds for primers 1( $\alpha\beta$ ) and 2( $\alpha$ )

59°C for 30 seconds for primers 3( $\alpha$ ) and 4( $\alpha$ )

57°C for 30 seconds for primers 1( $\alpha\beta$ ) and 2( $\beta$ )

63°C for 30 seconds for primers 3( $\beta$ ) and 4( $\beta$ )

extension at 75°C for 75 seconds.

The PCR products were confirmed to be the expected size by electrophoresis in a 2% agarose gel and ethidium bromide staining. The fragments were then purified by passage over a Wizard column (Promega) in accordance with the column manufacturer's recommendations.

The final coding sequence for TBP1(20-161)-hCG $\alpha$  was assembled by fusion PCR using primer 1( $\alpha\beta$ ) and primer 4( $\alpha$ ), and using as template the purified products from the p55 and hCG  $\alpha$  fragments obtained from the first PCR reactions. First the two templates, which due to the overlap between primers 2( $\alpha$ ) and 3( $\alpha$ ) could be denatured and annealed together, were passed through 10 cycles of PCR in the absence of any added primers. The conditions for these cycles were essentially the same as those used earlier, except that the annealing was done at 67°C and the extension was performed for 2 minutes. At the

end of these 10 cycles, primers 1( $\alpha\beta$ ) and 4( $\alpha$ ) were added, and another 10 cycles were performed. The conditions for this final set of reactions was the same as used earlier, except that an annealing temperature of 59°C was used, and the extension was performed for 75 seconds.

Analysis of the products of this reaction by electrophoresis in a 1% agarose gel confirmed that the expected fragment of about 1100bp was obtained. The reaction was passed over a Wizard column to purify the fragment, which was then digested with XbaI and BamHI and re-purified in a 0.7% low-melting point agarose gel. The purified fragment was subcloned into plasmid pSVL (Pharmacia), which had first been digested with XbaI and BamHI and gel purified on a 0.8% low-melting point agarose gel. Following ligation with T4 ligase, the mixture was used to transform AG1 *E. coli* and then plated onto LB/ampicillin plates for overnight culture at 37°C. Plasmid DNAs from ampicillin-resistant colonies were analyzed by digestion with XhoI and BamHI to confirm the presence of the insert (which is excised in this digest). Six clones were found to contain inserts, and one (clone 7) was selected for further advancement and designated pSVLTBPhCG $\alpha$  (containing TBP1(20-161)-hCG $\alpha$ ). Dideoxy DNA sequencing (using Sequenase™, U.S. Biochemicals, Cleveland, Ohio) of the insert in this vector confirmed that the construct was correct, and that no undesired changes had been introduced.

The final coding sequence for TBP1(20-161)-hCG $\beta$  was assembled in a manner similar to that described for TBP1(20-161)-hCG $\alpha$  using fusion PCR and primers 1( $\alpha\beta$ ) and 4( $\beta$ ), and using as template the purified products from the p55 and hCG  $\beta$  fragments obtained from the first PCR reactions. The resulting pSVL plasmid containing the insert of interest was designated pSVLTBPhCG $\beta$ .

b. TBP(20-190)-hCG

A second set of TBP-hCG proteins was prepared by modification of the TBP(20-161)-hCG constructs to produce an analog containing TBP spanning from Asp20 to Thr190, in place of the 20-161 region in the initial analog. This was done by

replacing the fragment between the BglII and XbaI sites in plasmid pSVLTBPhCG $\alpha$  with a PCR fragment containing the change. This PCR fragment was generated using fusion PCR. The primers were:

primer 1 . TTT TAG ATC TCT TCT TGC ACA GTG GAC  
(SEQ ID NO:18)

primer 2 TGT GGT GCC TGA GTC CTC AGT (SEQ ID  
NO:19)

primer 3 ACT GAG GAC TCA GGC ACC ACA GCC GGT GCT  
GCC CCA GGT TG (SEQ ID NO:20)

primer 4 TTT TTC TAG AGA AGC AGC AGC AGC CCA TG  
(SEQ ID NO:21)

Primers 1 and 2 were used to generate the sequence coding the additional p55 residues from 161-190. The PCR reaction was performed essentially as described earlier, using 1  $\mu$ g of each primer and pUC-p55 as template. Similarly, primers 3 and 4 were used to generate by PCR the linker between the 3'-end of the TBP-coding region, and the 5'-end of the hCG  $\alpha$  subunit coding region, using as a template plasmid pSVLTBPhCG $\alpha$ . Products from these PCR reactions were confirmed to be the correct size (about 296 bp and 121 bp respectively) by polyacrylamide gel electrophoresis (PAGE) on an 8% gel, and were then purified using a Wizard column. The design of primers 2 and 3 was such that they contained a region of overlap, so that the two PCR products (from primers 1 and 2, and from primers 3 and 4) could be annealed for fusion PCR with primers 1 and 4. Subsequent to the fusion reaction, the desired product of about 400 bp was confirmed and purified using a 1.5% agarose gel and a Wizard column. This DNA was then digested with BglII and XbaI, and ligated with BglII/XbaI-digested pSVLTBPhCG $\alpha$ . The presence of an insert in plasmids isolated from transformed AG1 *E. coli* was confirmed by digestion with BglII and XbaI. The new construct was designated pSVLTBP(20-190)-hCG $\alpha$ .

Similarly, plasmid pSVLTBPhCG $\beta$  was modified by substitution of the BglII-XcmI fragment. However, this was done by subcloning of a single PCR product, rather than with a

fusion PCR product. Primers 1 and 2b (see below) were used with pUC-p55 as the template.

primer 2b    TTT TCC ACA GCC AGG GTG GCA TTG ATG GGG  
                 CGG CAC CGT GGA CCA GCA CCA GCT GTG GTG  
                 CCT GAG TCC TCA GTG    (SEQ ID NO:22)

The resulting PCR product (about 337bp) was confirmed and purified as described above, digested with BglII and XcmI, and then ligated into BglII/XbaI-digested pSVLTBPhCG $\beta$ . The presence of an insert in plasmids isolated from transformed AG1 *E. coli* was confirmed by digestion with BglII and XcmI. The new construct was designated pSVLTBP(20-190)-hCG $\beta$ .

The new constructs were subsequently confirmed by DNA sequencing.

In addition to producing these new pSVL-based plasmids, these constructs were also subcloned into other expression vectors likely to be more suitable for stable expression in CHO, particularly vector D $\alpha$ , previously described as plasmid CLH3AXSV2DHFR (45). This was accomplished by converting a BamHI site flanking the inserts in the pSVL-based vectors to an XhoI site, and then excising the insert with XhoI and cloning it into XhoI digested D $\alpha$ .

## 2. Transient and stable expression of the hybrid proteins

Transfections of COS-7 cells (ATCC CRL 1651, ref. 46) for transient expression of the TBP-hCG hybrid proteins were performed using electroporation (47). Exponentially growing COS-7 cells were removed by trypsinization, collected by gentle centrifugation (800 rpm, 4 minutes), washed with cold phosphate buffered saline (PBS), pH 7.3-7.4, and then repelleted by centrifugation. Cells were resuspended at a concentration of  $5 \times 10^6$  cells per 400  $\mu$ l cold PBS and mixed with 10  $\mu$ g of plasmid DNA in a prechilled 2 mm gap electroporation cuvette. For cotransfections, 5  $\mu$ g of each plasmid were used. The cuvette and cells were chilled on ice for a further 10 minutes, and then subjected to electroporation using a BTX Model 600 instrument and conditions of 125 V, 950 $\mu$ F and R=8. Afterward the cells were set to cool on ice for 10 minutes,

transferred to a 15 ml conical tube containing 9.5 ml complete medium (Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 1% L-glutamine) at room temperature, and left at room temperature for 5 minutes. After gentle mixing in the 15 ml tube, the entire contents was seeded onto two P100 plates and placed into a 37°C, 5% CO<sub>2</sub> incubator. After 18 hours the media was changed, and in some cases the new media contained only 1% or 0% FBS. After another 72 hours, the conditioned media was harvested, centrifuged to remove cells, and then stored frozen at -70°C.

Transfections of CHO-DUKX (CHO) cells for transient or stable expression were performed using calcium phosphate precipitation of DNA. Twenty-four hours prior to the transfection, exponentially growing CHO cells were plated onto 100 mm culture plates at a density of  $7.5 \times 10^5$  cells per plate. On the day of the transfection, 10  $\mu$ g of plasmid DNA was brought to 0.5 ml in transfection buffer (see below), 31  $\mu$ l of 2 M CaCl<sub>2</sub> were added, the DNA-CaCl<sub>2</sub> solution was mixed by vortexing, and left to stand at room temperature for 45 minutes. After this the media was aspirated from the plates, the DNA was added to the cells using a sterile plastic pipette, and the cells were left at room temperature for 20 minutes. At the end of this period, 5 ml of complete  $\alpha$ (+)MEM containing 10% FBS was added to the plates, which were incubated at 37°C for 4-6 hours. The media was then aspirated off the plates, and the cells were subjected to a glycerol shock by incubating them with a solution of 15% glycerol in transfection buffer at 37°C for 3.5 minutes. After removal of the glycerol solution, the cells were washed twice with PBS, refed with 10 ml complete  $\alpha$ (+)MEM, 10% FBS, and returned to the 37°C incubator. For stable transfections, after 48 hours the cells were split 1:10 and fed with selection medium (complete  $\alpha$ -minus MEM (lacking nucleosides), 10% dialyzed FBS, and 0.02  $\mu$ M methotrexate). Non-transfected (non-resistant) cells were typically eliminated in 3-4 weeks, leaving a population of transfected, methotrexate-resistant cells.

### 3. Quantitation of expression

Secretion of the hybrid proteins by transfected cells was assessed using a commercial assay kit for soluble p55 (R&D Systems; Minneapolis, Minnesota) in accordance with the manufacturer's instructions. This assay also provides an estimate of the hybrid protein levels in conditioned and processed media, which served as the basis for selecting doses to be used in the bioassay.

### 4. Assessment of heterodimer formation

To assess the ability of the TBP-hCG subunit fusions to combine and form heterodimers, a sandwich immunoassay using antibodies to the hCG subunits was performed. In this assay, a monoclonal antibody to the hCG  $\beta$  subunit is coated onto microtiter plates and used for analyte capture. The primary detection antibody is a goat polyclonal raised against the human TSH  $\alpha$  subunit (#082422G - Biodesign International; Kennebunkport, Maine), which is in turn detected using a horse radish peroxidase conjugated rabbit anti-goat polyclonal antibody (Cappel; Durham, North Carolina).

Several different anti-hCG  $\beta$  subunit antibodies were used in this work, all of which show no detectable cross-reactivity with the free  $\alpha$  subunit. One of these antibodies (3/6) is used in the commercially available MAIAclone hCG assay kit (Biodata; Rome, Italy).

High-protein binding microtiter plates (Costar #3590) were coated with capture antibody by incubation (2 hours at 37°C) with 100  $\mu$ l/well of a 5  $\mu$ g/ml solution of antibody in coating buffer (PBS, pH 7.4, 0.1 mM  $\text{Ca}^{++}$ , 0.1 mM  $\text{Mg}^{++}$ ). After washing once with wash solution (PBS, pH 7.4 + 0.1% Tween 20) the plate is blocked by completely filling the wells ( $\approx$ 400  $\mu$ l/well) with blocking solution (3% bovine serum albumin (BSA; fraction V - A-4503 Sigma) in PBS, pH 7.4) and incubating for one hour at 37°C or overnight at 4°C. The plate is then washed twice with wash solution, and the reference and experimental samples, diluted in diluent (5 mg/ml BSA in PBS, pH 7.4) to yield a 100  $\mu$ l volume, are added. After incubating the samples and the plate for two hours at 37°C, the plate is again twice

washed with wash solution. The primary detection antibody, diluted 1:5000 in diluent, is added (100  $\mu$ l/well) and incubated for one hour at 37°C. The secondary detection antibody (HRP conjugated rabbit anti-goat Ig), diluted 1:5000 in diluent, is added (100  $\mu$ l/well) and after incubation for one hour at 37°C, the plate is washed three times with wash solution. One hundred  $\mu$ l of TMB substrate solution (Kirkegaard and Perry Laboratories) is added, the plate is incubated 20 minutes in the dark at room temperature, and then the enzymatic reaction is stopped by addition of 50  $\mu$ l/well 0.3M H<sub>2</sub>SO<sub>4</sub>. The plate is then analyzed using a microtiter plate reader set for a wavelength of 450 nm.

#### 5. Partial purification

To better quantitate the activities of these hybrid proteins, TBP-hCG hybrid proteins were partially purified by immunoaffinity chromatography. The antibody used was a monoclonal commercially available from R&D Systems (MAB #225). The column was CNBr-activated sepharose, charged with the antibody by following the manufacturer's (Pharmacia) instructions.

Conditioned media was collected from confluent T-175 flasks of each line using daily harvests of 50 ml SFMII media (GIBCO), five harvests for each line. The collections were subjected to centrifugation (1000 RPM) to remove cellular debris. The material was then assayed for TBP content using the commercial immunoassay and concentrated (Centricon units by Amicon; Beverly, Massachusetts) so that the apparent TBP concentration was about 50 ng/ml.

Ten ml of the concentrated TBP-hCG (sample #18873) was brought to approximately 1 M NaCl by addition of NaCl and adjustment of the solution to a conductivity of approximately 85 mS/cm. This was passed through a 0.5 ml anti-TBP immunoaffinity column. The flow-through was collected and run through the column a second time. After this the column was washed with 1 M NaCl in PBS. The bound TBP(20-161)-hCG was collected after elution with 50 mM citric acid (pH 2.5). The eluate (approximately 7 ml) was concentrated by filtration

using Amicon Centricon-10's in accordance with the manufacturer's (Amicon) instructions, to a volume of approximately 200  $\mu$ l. Approximately 800  $\mu$ l of PBS was added to bring the sample volume to 1 ml, which was stored at 4°C until tested by bioassay.

#### 6. Assessment of anti-TNF activity

Numerous *in vitro* TNF-induced cytotoxicity assays have been described for evaluating analogs of soluble TNF receptors. We utilized an assay employing a human breast carcinoma cell line, BT-20 cells (ATCC HTB 19). The use of these cells as the basis for a TNF bioassay has been described previously (48). These cells are cultured at 37°C in RPMI 1640 media supplemented with 10% heat-inactivated FBS. The cells were grown to a maximum 80-90% confluence, which entailed splitting every 3-4 days with a seeding density of about  $3 \times 10^6$  cells per T175cm<sup>2</sup> flask.

The BT-20 assay uses the inclusion of a cellular stain, crystal violet, as a detection method to assess survival of cells after treatment with TNF. Dead cells are unable to take up and retain the dye.

In brief, the protocol used for the assay of anti-TNF activity is the following. Recombinant human TNF $\alpha$  (R&D Systems) and the experimental samples are constituted in media (RPMI 1640 with 5% heat-inactivated FBS) and added to the wells of 96-well culture plates. The cells are then plated into these wells at a density of  $1 \times 10^5$  cells/well. The quantity of TNF $\alpha$  added was determined earlier in titration studies, and represents a dose at which about 50% of the cells are killed.

After addition of the samples, the cells are cultured for 48 hours at 39°C, after which the proportion of live cells is determined using crystal violet staining and a microtiter plate reader (570 nm).

### RESULTS

#### 1. Constructs under study

The designs of the hybrid proteins studied are briefly summarized below; two control proteins, a monomeric



soluble p55 (r-hTBP-1) and a dimeric TBP-immunoglobulin fusion protein (TBP-IgG3) (prepared essentially as described in (10)), were studied for comparative purposes.

<u>Construct</u>	<u>TBP N-term</u>	<u>TBP C-term</u>	<u>Fusion partner</u>
r-hTBP-1	mix of 9 and 20	180	none
TBP-IgG3	mix of 9 and 20	190	IgG3 heavy chain constant region
TBP(20-161)-hCG	20	161	hCG $\alpha$ and hCG $\beta$ (heterodimer)
TBP(20-190)-hCG	20	190	hCG $\alpha$ and hCG $\beta$ (heterodimer)

The sequences of the DNAs encoding, TBP(20-190)-hCG and TBP(20-161)-hCG are provided in Figures 1 and 2, respectively. A schematic summary of the constructs is provided in Figure 3.

## 2. Secretion of TBP-hCG proteins

All of the constructs tested were found to be produced and secreted into culture media by transfected mammalian cells. Data illustrating this are shown in Tables 1 and 2.

## 3. TBP-hCG( $\alpha/\beta$ ) fusion proteins assemble into heterodimers

The combination of TBP-hCG $\alpha$  and TBP-hCG $\beta$  was confirmed using the sandwich assay for the hCG heterodimer. Only the combined transfection of  $\alpha$  and  $\beta$  subunit fusions resulted in heterodimer detection (Table 3).

## 4. TBP-hCG hybrid proteins exhibit increased activity over TBP monomer

Hybrid proteins produced in either COS-7 or CHO cells were found to be potent inhibitors of TNF $\alpha$  in the BT-20 bioassay. Some of the samples tested are summarized in Table 4.

Negative controls (conditioned media from mock transfections) were included for the 1x media samples.

As illustrated in Figures 4-6 (points on y-axis), addition of TNF (2.5 ng/ml) results in a clear reduction in live cell number (as assessed by OD 570). In every case, active samples have as a maximal protective effect the restoration of cell viability to the level seen in the absence of added TNF (i.e., the control labeled "cells alone").

The positive controls, r-hTBP-1 and TBP-IgG3, are both protective, showing a clear dose-dependence and ED50s of approximately 100 ng/ml for the r-hTBP-1 (Figs. 4-6) and about 1.5 ng/ml for TBP-IgG3 (Fig. 4) respectively.

The TBP-hCG constructs from 1x media (CHO or COS) or from the immunopurification show dose-dependent protection, with approximate ED50s ranging from 2-11 ng/ml (Figs. 4-6).

The results from the *in vitro* bioassay are reported in Table 5. The data indicate that the hybrid proteins inhibit TNF cytotoxicity, and that they are substantially more potent than the TBP monomer. The negative controls were devoid of protective activity.

In addition to the possibility that dimerization of TBP may increase potency, it is also possible that the activity of the hybrid proteins are not related to dimeric interaction with TBP, but rather to steric inhibition due to the partner of the hybrid interfering with soluble TBP/TNF binding to cell-surface TNF receptors.

All references cited herein, including journal articles or abstracts, published or corresponding U.S. or foreign patent applications, issued U.S. or foreign patents, or any other references, are entirely incorporated by reference herein, including all data, tables, figures, and text presented in the cited references. Additionally, the entire contents of the references cited within the references cited herein are also entirely incorporated by reference.

Reference to known method steps, conventional method steps, known methods or conventional methods is not in any way an admission that any aspect, description or embodiment of the present invention is disclosed, taught or suggested in the relevant art.

The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying knowledge within the skill of the art (including the contents of the references cited herein), readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance presented herein, in combination with the knowledge of one of ordinary skill in the art.

## TABLES

Table 1: COS-7 transient expression (TBP ELISA)	
Hybrid Protein	Concentration (pg/ml)
TBP1	66
TBP-hCG $\alpha$ (20-161)	5.1
TBP-hCG $\beta$ (20-161)	0.5
TBP-hCG(20-161)	2.7
control	<0.25

Constructs were expressed using pSVL (Pharmacia)

Table 2: COS-7 transient expression (TBP ELISA)	
Hybrid protein	Concentration (ng/ml)
TBP1	131
TBP-hCG $\alpha$ (20-190)	81
TBP-hCG $\beta$ (20-190)	9
TBP-hCG(20-190)	62
control	<1

Constructs were expressed using a mouse metallothionein promoter-containing vector - pD $\alpha$

Table 3: COS-7 transient expression (hCG heterodimer assay)	
Hybrid Protein	Concentration (ng/ml)
TBP1	<0.2
TBP-hCG $\alpha$ (20-190)	<0.2
TBP-hCG $\beta$ (20-190)	<0.2
TBP-hCG(20-190)	38
control	<0.2

Constructs were expressed using a mouse  
metallothionein promoter-containing vector - pD $\alpha$

Table 4: Samples tested for anti-TNF activity		
<i>Construct</i>	<i>Cell source</i>	<i>Nature of sample</i>
r-hTBP-1	CHO	purified
TBP-IgG3	CHO	1x conditioned media
TBP(20-161)-hCG	CHO	immunopurified (anti-TBP)
TBP(20-190)-hCG	CHO	1x conditioned media
TBP(20-190)-hCG	COS	1x conditioned media

Table 5 :Preliminary Assessment of the hybrid proteins in TNF Cytotoxicity Assay		
<i>Construct</i>	<i>Fusion partner</i>	<i>Anti-TNF activity (ED50) in BT-20 bioassay*</i>
r-hTBP-1	none	100 ng/ml
TBP-IgG3	IgG3 heavy chain constant region	1.5 ng/ml
TBP(20-161)-hCG	hCG $\alpha$ and hCG $\beta$ (heterodimer)	2 ng/ml
TBP(20-190)-hCG	hCG $\alpha$ and hCG $\beta$ (heterodimer)	8-11 ng/ml

\*\*The quantitation of material for dosing and estimation of ED50 was made using the TBP ELISA.

## REFERENCES

1. Smith, R.A. et al., J. Biol. Chem. 262:6951-6954, 1987.
2. Eck, M.J. et al., J. Biol. Chem. 264:17595-17605, 1989.
3. Jones, E.Y. et al., Nature 338:225-228, 1989.
4. Eck, M.J. et al., J. Biol. Chem. 267:2119-2122, 1992.
5. Pierce, J.G. et al., Annu. Rev. Biochem. 50:465-495, 1981.
6. Lapthorn, A.J. et al., Nature 369:455-461, 1994.
7. Wu, H., et al., Structure 2:545-550, 1994.
8. Engelmann, H., et al., J. Biol. Chem. 265:14497-14504, 1990.
9. Adam, D. et al., J. Biol. Chem. 270:17482-17487, 1995.
10. Loetscher, H.R., et al., J. Biol. Chem. 266:18324-18329, 1991.
11. Banner, D.W., et al., Cell 73:431-445, 1993.
12. Pennica, D., et al., Biochemistry 32:3131-3138, 1993.
13. Engelmann, H. et al., J. Biol. Chem. 265:1531-1536, 1990.
14. Van Zee, K.J. et al., Proc. Natl. Acad. Sci. USA 89:4845-4849, 1992.
15. Aderka, D. et al., J. Exp. Med. 175:323-329, 1992.
16. Mohler, K.M., et al., J. Immunol. 151:1548-1561, 1993.
17. Bertini, R., et al., Eur. Cytokine Netw., 1993.
18. Piguet, P.F., et al., Immunology 77:510-514, 1992.
19. Williams, R.O., et al., Immunology 84:433-439, 1995.
20. Capon, D.J., et al., Nature 337: 525-531, 1989.

21. Ashkenazi, A., et al., Proc. Natl. Acad. Sci. 88:10535-10539, 1991.
22. Suitters, A.J., et al. J. Exp. Med. 179:849-856, 1994.
23. Nolan, O. et al., Biochim. Biophys. Acta 1040:1-11, 1990.
24. Rodrigues, M.L., et al., J. Immunol. 151:6954-6961, 1993.
25. Chang, H.-C., et al., Proc. Natl. Acad. Sci. USA 91:11408-11412, 1994.
26. Wu, Z., et al., J. Biol. Chem. 270:16039-16044, 1995.
27. Bazzoni, F. et al, Proc. Natl. Acad. Sci. USA 92:5376-5380, 1995.
28. Boldin, M.P., et al., J. Biol. Chem. 270:387-391, 1995.
29. Vu, T.-K.H., et al., Cell, 64:1057-1068, 1991.
30. Song, H.Y., et al., J. Biol. Chem. 269:22492-22495, 1994.
31. Russell, D.A., et al., J. Infectious Diseases 171:1528-1538, 1995.
32. Rao C.V. et al., Am. J. Obstet. Gynecol., 146, 65-68, 1983.
33. Damewood M.D. et al., Fertil. Steril. 52, 398-400, 1989.
34. Chen, F., et al., Mol. Endocrinol. 6:914-919, 1992.
35. Bielinska, M., et al., J. Cell Biol. 111:330a, 1990.
36. Furuhashi, M., et al., Mol Endocrinol. 9:54-63, 1995.
37. Sugahara, T., et al., Proc. Natl. Acad. Sci. USA 92:2041-2045, 1995.
38. Johnson, G.A., et al., Biol. Reprod. 52:68-73, 1995.
39. Urlaub, G. and Chasin, L. Proc. Natl. Acad. Sci. USA 77:4216-4220, 1980.



40. Nophar, Y., et al., EMBO J. 9:3269-3278, 1990.
41. Fiddes, J.C. et al., Nature 281:351-356, 1979.
42. Fiddes, J.C. et al., Nature 286:684-687, 1980.
43. Elion, E.A., in Current Protocols in Molecular Biology, eds. Ausuble, FM. et al., John Wiley & Sons, 1993.
44. Campbell, R., Proc. Natl. Acad. Sci. USA 88:760-764, 1991.
45. Cole E.S. et al., Biotechnology, 11, 1014-1024, 1993.
46. Gluzman, Y., Cell 23:175-182, 1981.
47. Chu, G. et al., Nucl. Acid Res. 15:1311-1326, 1987.
48. Yen, J. et al., J. Immunotherapy 10:174-181, 1991.

# SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Campbell, Robert K.  
Jameson, Bradford A.  
Chappel, Scott C.
- (ii) TITLE OF INVENTION: HYBRID PROTEINS
- (iii) NUMBER OF SEQUENCES: 22
- (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: BROWDY AND NEIMARK  
(B) STREET: 419 Seventh Street N.W., Ste. 300  
(C) CITY: Washington  
(D) STATE: D.C.  
(E) COUNTRY: USA  
(F) ZIP: 22207
- (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Floppy disk  
(B) COMPUTER: IBM PC compatible  
(C) OPERATING SYSTEM: PC-DOS/MS-DOS  
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: 60/011,936  
(B) FILING DATE: 20 February 1996  
(C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: Browdy, Roger L.  
(B) REGISTRATION NUMBER: 25,618  
(C) REFERENCE/DOCKET NUMBER: CAMPBELL=2A
- (ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: (202) 628-5197  
(B) TELEFAX: (202) 737-3528

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 1049 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 278..1047

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCCACATGGC TACAGGTAAG CGCCCCTAAA ATCCCTTTGG GCACAATGTG TCCTGAGGGG	60
AGAGGCAGCG ACCTGTAGAT GGGACGGGGG CACTAACCT CAGGTTTGGG GCTTCTCAAT	120
CTCACTATCG CCATGTAAGC CCAGTATTTG GCCAATCTCA GAAAGCTCCT CCTCCCTGGA	180
GGGATGGAGA GAGAAAAACA AACAGCTCCT GGAGCAGGGA GAGTGCTGGC CTCTTGCTCT	240
CCGGCTCCCT CTGTTGCCCT CTGGTTTCTC CCCAGGC TCC CGG ACG TCC CTG CTC	295
Ser Arg Thr Ser Leu Leu	
1 5	

CTG GCT TTT GGC CTG CTC TGC CTG CCC TGG CTT CAA GAG GGC AGT GCC	343
Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp Leu Gln Glu Gly Ser Ala	
10 15 20	
GAT AGT GTG TGT CCC CAA GGA AAA TAT ATC CAC CCT CAA AAT AAT TCC	391
Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser	
25 30 35	
ATT TGC TGT ACC AAG TGC CAC AAA GGA ACC TAC TTG TAC AAT GAC TGT	439
Ile Cys Cys Thr Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys	
40 45 50	
CCA GGC CCG GGG CAG GAT ACG GAC TGC AGG GAG TGT GAG AGC GGC TCC	487
Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser	
55 60 65 70	
TTC ACC GCT TCA GAA AAC CAC CTC AGA CAC TGC CTC AGC TGC TCC AAA	535
Phe Thr Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys	
75 80 85	
TGC CGA AAG GAA ATG GGT CAG GTG GAG ATC TCT TCT TGC ACA GTG GAC	583
Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp	
90 95 100	
CGG GAC ACC GTG TGT GGC TGC AGG AAG AAC CAG TAC CGG CAT TAT TGG	631
Arg Asp Thr Val Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr Trp	
105 110 115	
AGT GAA AAC CTT TTC CAG TGC TTC AAT TGC AGC CTC TGC CTC AAT GGG	679
Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn Gly	
120 125 130	
ACC GTG CAC CTC TCC TGC CAG GAG AAA CAG AAC ACC GTG TGC ACC TGC	727
Thr Val His Leu Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr Cys	
135 140 145 150	
CAT GCA GGT TTC TTT CTA AGA GAA AAC GAG TGT GTC TCC TGT GCC GGT	775
His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ala Gly	
155 160 165	
GCT GCC CCA GGT TGC CCA GAA TGC ACG CTA CAG GAA AAC CCA TTC TTC	823
Ala Ala Pro Gly Cys Pro Glu Cys Thr Leu Gln Glu Asn Pro Phe Phe	
170 175 180	
TCC CAG CCG GGT GCC CCA ATA CTT CAG TGC ATG GGC TGC TGC TTC TCT	871
Ser Gln Pro Gly Ala Pro Ile Leu Gln Cys Met Gly Cys Cys Phe Ser	
185 190 195	
AGA GCA TAT CCC ACT CCA CTA AGG TCC AAG AAG ACG ATG TTG GTC CAA	919
Arg Ala Tyr Pro Thr Pro Leu Arg Ser Lys Lys Thr Met Leu Val Gln	
200 205 210	
AAG AAC GTC ACC TCA GAG TCC ACT TGC TGT GTA GCT AAA TCA TAT AAC	967
Lys Asn Val Thr Ser Glu Ser Thr Cys Cys Val Ala Lys Ser Tyr Asn	
215 220 225 230	
AGG GTC ACA GTC ATG GGG GGT TTC AAA GTG GAG AAC CAC ACG GGG TGC	1015
Arg Val Thr Val Met Gly Gly Phe Lys Val Glu Asn His Thr Gly Cys	
235 240 245	
CAC TGC AGT ACT TGT TAT TAT CAC AAA TCT TA AG	1049
His Cys Ser Thr Cys Tyr Tyr His Lys Ser	
250 255	

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 256 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Ser Arg Thr Ser Leu Leu Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp  
1 5 10 15  
Leu Gln Glu Gly Ser Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile  
20 25 30  
His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys Gly Thr  
35 40 45  
Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg  
50 55 60  
Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu Arg His  
65 70 75 80  
Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile  
85 90 95  
Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg Lys Asn  
100 105 110  
Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys  
115 120 125  
Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu Lys Gln  
130 135 140  
Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu  
145 150 155 160  
Cys Val Ser Cys Ala Gly Ala Ala Pro Gly Cys Pro Glu Cys Thr Leu  
165 170 175  
Gln Glu Asn Pro Phe Phe Ser Gln Pro Gly Ala Pro Ile Leu Gln Cys  
180 185 190  
Met Gly Cys Cys Phe Ser Arg Ala Tyr Pro Thr Pro Leu Arg Ser Lys  
195 200 205  
Lys Thr Met Leu Val Gln Lys Asn Val Thr Ser Glu Ser Thr Cys Cys  
210 215 220  
Val Ala Lys Ser Tyr Asn Arg Val Thr Val Met Gly Gly Phe Lys Val  
225 230 235 240  
Glu Asn His Thr Gly Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser  
245 250 255

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1202 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 279..1199

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CTCGAGATGG CTACAGGTAA GCGCCCCTAA AATCCCTTTG GGCACAATGT GTCCTGAGGG	60
GAGAGGTAGC GACCTGTAGA TGGGACGGGG GCACTAACCC TGAGGTTTGG GGCTTCTGAA	120
TGTGAGTATC GCCATGTAAG CCCAGTATTT GGCCAATGTC AGAAAGCTCC TGGTCCCTGG	180
AGGGATGGAG AGAGAAAAAC AAACAGCTCC TGGAGCAGGG AGAGTGCTGG CCTCTTGCTC	240
TCCGGCTCCC TCTGTTGCCC TGTGGTTTCT CCCCAGGC TCC CGG ACG TCC CTG	293
Ser Arg Thr Ser Leu 260	
CTC CTG GCT TTT GGC CTG CTC TGC CTG CCC TGG CTT CAA GAG GGC AGT	341
Leu Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp Leu Gln Glu Gly Ser	
265 270 275	
GCC GAT AGT GTG TGT CCC CAA GGA AAA TAT ATC CAC CCT CAA AAT AAT	389
Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn	
280 285 290	
TCG ATT TGC TGT ACC AAG TGC CAC AAA GGA ACC TAC TTG TAC AAT GAC	437
Ser Ile Cys Cys Thr Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp	
295 300 305	
TGT CCA GGC CCG GGG CAG GAT ACG GAC TGC AGG GAG TGT GAG AGC GGC	485
Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly	
310 315 320 325	
TCT TTC ACC GCT TCA GAA AAC CAC CTC AGA CAC TGC CTC AGC TGC TCC	533
Ser Phe Thr Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser	
330 335 340	
AAA TGC CGA AAG GAA ATG GGT CAG GTG GAG ATC TCT TCT TGC ACA GTG	581
Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val	
345 350 355	
GAC CGG GAC ACC GTG TGT GGC TGC AGG AAG AAC CAG TAC CGG CAT TAT	629
Asp Arg Asp Thr Val Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr	
360 365 370	
TGG AGT GAA AAC CTT TTC CAG TGC TTC AAT TGC AGC CTC TGC CTC AAT	677
Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn	
375 380 385	
GGG ACC GTG CAC CTC TCC TGC CAG GAG AAA CAG AAC ACC GTG TGC ACC	725
Gly Thr Val His Leu Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr	
390 395 400 405	
TGC CAT GCA GGT TTC TTT CTA AGA GAA AAC GAG TGT GTC TCC TGT GCT	773
Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ala	
410 415 420	
GGT GCT GGT CCA CGG TGC CGC CCC ATC AAT GCC ACC CTG GCT GTG GAG	821
Gly Ala Gly Pro Arg Cys Arg Pro Ile Asn Ala Thr Leu Ala Val Glu	
425 430 435	
AAG GAG GGC TGC CCC GTG TGC ATC ACC GTC AAC ACC ACC ATC TGT GCC	869
Lys Glu Gly Cys Pro Val Cys Ile Thr Val Asn Thr Thr Ile Cys Ala	
440 445 450	

GGC TAC TGC CCC ACC ATG ACC CGC GTG CTG CAG GGG GTC CTC CCC GCC	917
Gly Tyr Cys Pro Thr Met Thr Arg Val Leu Gln Gly Val Leu Pro Ala	
455 460 465	
CTG CCT CAG GTG GTG TGC AAC TAC CGC GAT GTG CGC TTC GAG TCC ATC	965
Leu Pro Gln Val Val Cys Asn Tyr Arg Asp Val Arg Phe Glu Ser Ile	
470 475 480 485	
CGG CTC CCT GGC TGC CCG CGC GGC GTG AAC CCC GTG GTC TCC TAC GCT	1013
Arg Leu Pro Gly Cys Pro Arg Gly Val Asn Pro Val Val Ser Tyr Ala	
490 495 500	
GTG GCT CTC AGC TGT CAA TGT GCA CTC TGC CGC CGC AGC ACC ACT GAC	1061
Val Ala Leu Ser Cys Gln Cys Ala Leu Cys Arg Arg Ser Thr Thr Asp	
505 510 515	
TGC GGG GGT CCC AAG GAC CAC CCC TTG ACC TGT GAT GAC CCC CGC TTC	1109
Cys Gly Gly Pro Lys Asp His Pro Leu Thr Cys Asp Asp Pro Arg Phe	
520 525 530	
CAG GAC TCC TCT TCC TCA AAG GCC CCT CCC CCC AGC CTT CCA AGC CCA	1157
Gln Asp Ser Ser Ser Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro	
535 540 545	
TCC CGA CTC CCG GGG CCC TCG GAC ACC CCG ATC CTC CCA CAA TAA	1202
Ser Arg Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln	
550 555 560	

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 307 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Ser	Arg	Thr	Ser	Leu	Leu	Leu	Ala	Phe	Gly	Leu	Leu	Cys	Leu	Pro	Trp
1				5					10					15	
Leu	Gln	Glu	Gly	Ser	Ala	Asp	Ser	Val	Cys	Pro	Gln	Gly	Lys	Tyr	Ile
			20					25					30		
His	Pro	Gln	Asn	Asn	Ser	Ile	Cys	Cys	Thr	Lys	Cys	His	Lys	Gly	Thr
			35				40					45			
Tyr	Leu	Tyr	Asn	Asp	Cys	Pro	Gly	Pro	Gly	Gln	Asp	Thr	Asp	Cys	Arg
	50				55						60				
Glu	Cys	Glu	Ser	Gly	Ser	Phe	Thr	Ala	Ser	Glu	Asn	His	Leu	Arg	His
	65				70					75					80
Cys	Leu	Ser	Cys	Ser	Lys	Cys	Arg	Lys	Glu	Met	Gly	Gln	Val	Glu	Ile
				85					90					95	
Ser	Ser	Cys	Thr	Val	Asp	Arg	Asp	Thr	Val	Cys	Gly	Cys	Arg	Lys	Asn
			100					105					110		
Gln	Tyr	Arg	His	Tyr	Trp	Ser	Glu	Asn	Leu	Phe	Gln	Cys	Phe	Asn	Cys
		115					120					125			
Ser	Leu	Cys	Leu	Asn	Gly	Thr	Val	His	Leu	Ser	Cys	Gln	Glu	Lys	Gln
	130					135					140				

Asn	Thr	Val	Cys	Thr	Cys	His	Ala	Gly	Phe	Phe	Leu	Arg	Glu	Asn	Glu	145	150	155	160
Cys	Val	Ser	Cys	Ala	Gly	Ala	Gly	Pro	Arg	Cys	Arg	Pro	Ile	Asn	Ala	165	170	175	
Thr	Leu	Ala	Val	Glu	Lys	Glu	Gly	Cys	Pro	Val	Cys	Ile	Thr	Val	Asn	180	185	190	
Thr	Thr	Ile	Cys	Ala	Gly	Tyr	Cys	Pro	Thr	Met	Thr	Arg	Val	Leu	Gln	195	200	205	
Gly	Val	Leu	Pro	Ala	Leu	Pro	Gln	Val	Val	Cys	Asn	Tyr	Arg	Asp	Val	210	215	220	
Arg	Phe	Glu	Ser	Ile	Arg	Leu	Pro	Gly	Cys	Pro	Arg	Gly	Val	Asn	Pro	225	230	235	240
Val	Val	Ser	Tyr	Ala	Val	Ala	Leu	Ser	Cys	Gln	Cys	Ala	Leu	Cys	Arg	245	250	255	
Arg	Ser	Thr	Thr	Asp	Cys	Gly	Gly	Pro	Lys	Asp	His	Pro	Leu	Thr	Cys	260	265	270	
Asp	Asp	Pro	Arg	Phe	Gln	Asp	Ser	Ser	Ser	Ser	Lys	Ala	Pro	Pro	Pro	275	280	285	
Ser	Leu	Pro	Ser	Pro	Ser	Arg	Leu	Pro	Gly	Pro	Ser	Asp	Thr	Pro	Ile	290	295	300	
Leu	Pro	Gln														305			

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1147 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 278..1132

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

TCGAGATGGC TACAGGTAAG CGCCCCTAAA ATCCCTTTGG GCACAATGTG TCCTGAGGGG	60
AGAGGCAGCG ACCTGTAGAT GGGACGGGGG CACTAACCCT CAGGTTTGGG GCTTTTGAAT	120
GTGAGTATGG CCATGTAAGC CCAGTATTTG CCCAATCTCA GAAAGCTCCT GTCCCTGGA	180
GGGATGGAGA GAGAAAAACA AACAGCTCCT GGAGCAGGGA CACTCCTGGC CTCTTGCTCT	240
GCGGCTCCGT GTGTTGCCCT GTGGTTTCTC CCCACGC TCC CGG ACG TCC CTG CTC	295
Ser Arg Thr Ser Leu Leu	310
CTG GCT TTT GGC CTG CTC TGC CTG CCC TGG CTT CAA GAG GGC AGT GCC	343
Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp Leu Gln Glu Gly Ser Ala	315 320 325

GAT Asp 330	AGT Ser	GTG Val	TGT Cys	CCC Pro	CAA Gln 335	GGA Gly	AAA Lys	TAT Tyr	ATC Ile	CAC His 340	CCT Pro	CAA Gln	AAT Asn	AAT Asn	TCG Ser 345	391
ATT Ile	TGC Cys	TGT Cys	ACC Thr	AAG Lys 350	TGC Cys	CAC His	AAA Lys	GGA Gly	ACC Thr 355	TAC Tyr	TTG Leu	TAC Tyr	AAT Asn	GAC Asp 360	TGT Cys	439
CCA Pro	GGC Gly	CCG Pro	GGG Glu 365	CAG Gln	GAT Asp	ACC Thr	GAC Asp	TGC Cys 370	AGG Arg	GAG Glu	TGT Cys	GAG Glu	AGC Ser 375	GGC Gly	TCC Ser	487
TTC Phe	ACC Thr	GCT Ala 380	TCA Ser	GAA Glu	AAC Asn	CAC His	CTC Leu 385	AGA Arg	CAC His	TGC Cys	CTC Leu 390	AGC Ser	TGC Cys	TCC Ser	AAA Lys	535
TGC Cys 395	CGA Arg	AAG Lys	GAA Glu	ATG Met	GGT Gly	CAG Gln 400	GTG Val	GAG Glu	ATC Ile	TCT Ser 405	TCT Ser	TGC Cys	ACA Thr	GTG Val	GAC Asp	583
CGG Arg 410	GAC Asp	ACC Thr	GTG Val	TGT Cys	GGC Gly 415	TGC Cys	AGG Arg	AAG Lys	AAC Asn	CAG Gln 420	TAC Tyr	CGG Arg	CAT His	TAT Tyr	TGG Trp 425	631
AGT Ser	GAA Glu	AAC Asn	CTT Leu	TTC Phe 430	CAG Gln	TGC Cys	TTC Phe	AAT Asn	TGC Cys 435	ACC Thr	CTC Leu	TGC Cys	CTC Leu	AAT Asn 440	GGG Gly	679
ACC Thr	GTG Val	CAC His	CTC Leu 445	TCC Ser	TGT Cys	CAG Gln	GAG Glu	AAA Lys 450	CAG Gln	AAC Asn	ACC Thr	GTC Val	TGC Cys 455	ACC Thr	TGC Cys	727
CAT His	GCA Ala	GGT Gly 460	TTC Phe	TTT Phe	CTA Leu	AGA Arg	GAA Glu 465	AAC Asn	GAG Glu	TGT Cys	GTC Val 470	TCC Ser	TGT Cys	AGT Ser	AAC Asn	775
TGT Cys 475	AAG Lys	AAA Lys	AGC Ser	CTG Leu	GAG Glu	TGC Cys 480	ACG Thr	AAG Lys	TTG Leu	TCC Ser	CTA Leu 485	CCC Pro	CAG Gln	ATT Ile	GAG Glu	823
AAT Asn 490	GTT Val	AAG Lys	GGC Gly	ACT Thr	GAG Glu 495	GAC Asp	TCA Ser	GGC Gly	ACC Thr	ACA Thr 500	GCC Ala	GGT Gly	GCT Ala	GCC Ala	CCA Pro 505	871
GGT Gly	TGC Cys	CCA Pro	GAA Glu	TGC Cys 510	ACG Thr	CTA Leu	CAG Gln	GAA Glu 515	AAC Asn	CCA Pro	TTC Phe	TTC Phe	TCC Ser	CAG Gln 520	CCG Pro	919
GGT Gly	GCC Ala	CCA Pro	ATA Ile 525	CTT Leu	CAG Gln	TGC Cys	ATG Met	GGC Gly 530	TGC Cys	TGC Cys	TTC Phe	TCT Ser	AGA Arg 535	GCA Ala	TAT Tyr	967
CCC Pro	ACT Thr	CCA Pro 540	CTA Leu	AGG Arg	TCC Ser	AAG Lys	AAG Lys 545	ACG Thr	ATG Met	TTG Leu	GTC Val	CAA Gln 550	AAG Lys	AAC Asn	GTC Val	1015
ACC Thr 555	TCA Ser	GAG Glu	TCC Ser	ACT Thr	TGC Cys	TGT Cys	GTA Val 560	GCT Ala	AAA Lys	TCA Ser	TAT Tyr 565	AAC Asn	AGG Arg	GTC Val	ACA Thr	1063
GTA Val 570	ATG Met	GGG Gly	GGT Gly	TTC Phe	AAA Lys 575	GTG Val	GAG Glu	AAC Asn	CAC His	ACG Thr 580	GCG Ala	TGC Cys	CAC His	TGC Cys	AGT Ser 585	1111
ACT Thr	TGT Cys	TAT Tyr	TAT Tyr	CAC His 590	AAA Lys	TCT Ser	TAAGGATCCC TCGAG									1147



(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 285 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ser Arg Thr Ser Leu Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp  
1 5 10 15  
Leu Gln Glu Gly Ser Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile  
20 25 30  
His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys Gly Thr  
35 40 45  
Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg  
50 55 60  
Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu Arg His  
65 70 75 80  
Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile  
85 90 95  
Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys Arg Lys Asn  
100 105 110  
Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys  
115 120 125  
Thr Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys Gln Glu Lys Gln  
130 135 140  
Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu  
145 150 155 160  
Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Leu Glu Cys Thr Lys Leu  
165 170 175  
Ser Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Ser Gly Thr  
180 185 190  
Thr Ala Gly Ala Ala Pro Gly Cys Pro Glu Cys Thr Leu Gln Glu Asn  
195 200 205  
Pro Phe Phe Ser Gln Pro Gly Ala Pro Ile Leu Gln Cys Met Gly Cys  
210 215 220  
Cys Phe Ser Arg Ala Tyr Pro Thr Pro Leu Arg Ser Lys Lys Thr Met  
225 230 235 240  
Leu Val Gln Lys Asn Val Thr Ser Glu Ser Thr Cys Cys Val Ala Lys  
245 250 255  
Ser Tyr Asn Arg Val Thr Val Met Gly Gly Phe Lys Val Glu Asn His  
260 265 270  
Thr Ala Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser  
275 280 285

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1301 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 279..1287

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CTCGAGATGG CTACAGGTAA GCGCCCCTAA AATCCCTTTG GGCACAATGT GTCCTGAGGG	60
GAGAGGCAGC GACCTGTAGA TGGGACGGGG GCACTAACCC TCAGGTTTGG GGCTTCTGAA	120
TGTGAGTATC GCCATGTAAG CCCAGTATTT GGCCAATGTC AGAAAGCTCC TGGTCCCTGG	180
AGGGATGGAG AGAGAAAAAC AAACACCTCC TGGAGCAGGG AGAGTGCTGC CCTCTTGCTC	240
TCCGGCTCCC TCTGTTGCCC TCTGGTTTCT CCCCAGGC TCC CGG ACG TCC CTG	293
	Ser Arg Thr Ser Leu
	290
CTC CTG GCT TTT GGC CTG CTC TGC CTG CCC TGG CTT CAA GAG GGC AGT	341
Leu Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp Leu Gln Glu Gly Ser	
	295 300 305
GCC GAT AGT GTG TGT CCC CAA GGA AAA TAT ATC CAC CCT CAA AAT AAT	389
Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn	
	310 315 320
TCG ATT TGC TGT ACC AAG TGC CAC AAA GGA ACC TAC TTG TAC AAT GAC	437
Ser Ile Cys Cys Thr Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp	
	325 330 335
TGT CCA GGC CCG GGG CAG GAT ACG GAC TGC AGG GAG TGT GAG AGC GGC	485
Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly	
	340 345 350
TCC TTC ACC GCT TCA GAA AAC CAC CTC AGA CAC TGC CTC AGC TGC TCC	533
Ser Phe Thr Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser	
	355 360 365 370
AAA TGC CGA AAG GAA ATG GGT CAG GTG GAG ATC TCT TCT TGC ACA GTG	581
Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val	
	375 380 385
GAC CGG GAC ACC GTG TGT GGC TGC AGG AAG AAC CAG TAC CGG CAT TAT	629
Asp Arg Asp Thr Val Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr	
	390 395 400
TGG AGT GAA AAC CTT TTC CAG TGC TTC AAT TGC AGC CTC TGC CTC AAT	677
Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn	
	405 410 415
GGG ACC GTG CAC CTC TCC TGC CAG GAG AAA CAG AAC ACC GTG TGC ACC	725
Gly Thr Val His Leu Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr	
	420 425 430
TGC CAT GCA GGT TTC TTT CTA AGA GAA AAC GAG TGT GTC TCC TGT AGT	773
Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ser	
	435 440 445 450

AAC TGT AAG AAA AGC CTG GAG TGC ACG AAG TTG TGC CTA CCC CAG ATT	821
Asn Cys Lys Lys Ser Leu Glu Cys Thr Lys Leu Cys Leu Pro Gln Ile	
455 460 465	
GAG AAT GTT AAG GGC ACT GAG GAC TCA GGC ACC ACA GCT GGT GCT GGT	869
Glu Asn Val Lys Gly Thr Glu Asp Ser Gly Thr Thr Ala Gly Ala Gly	
470 475 480	
CCA CGG TGC CGC CCC ATC AAT GCC ACC CTG GCT GTG GAG AAG GAG GGC	917
Pro Arg Cys Arg Pro Ile Asn Ala Thr Leu Ala Val Glu Lys Glu Gly	
485 490 495	
TGC CCC GTG TGC ATC ACC GTC AAC ACC ACC ATC TGT GCC GGC TAC TGC	965
Cys Pro Val Cys Ile Thr Val Asn Thr Thr Ile Cys Ala Gly Tyr Cys	
500 505 510	
CCC ACC ATG ACC CGC GTG CTG CAG GGG GTC CTG CCG GCC CTG CCT CAG	1013
Pro Thr Met Thr Arg Val Leu Gln Gly Val Leu Pro Ala Leu Pro Gln	
515 520 525 530	
GTG GTG TGC AAC TAC CGC GAT GTG CGC TTC GAG TCC ATC CGG CTC CCT	1061
Val Val Cys Asn Tyr Arg Asp Val Arg Phe Glu Ser Ile Arg Leu Pro	
535 540 545	
GGC TGC CCG CGC GGC GTG AAC CCC GTG GTC TCC TAC GCC GTG GCT CTC	1109
Gly Cys Pro Arg Gly Val Asn Pro Val Val Ser Tyr Ala Val Ala Leu	
550 555 560	
AGC TGT CAA TGT GCA CTC TGC CGC CGC AGC ACC ACT GAC TGC GGG GGT	1157
Ser Cys Gln Cys Ala Leu Cys Arg Arg Ser Thr Thr Asp Cys Gly Gly	
565 570 575	
CCC AAG GAC CAC CCC TTG ACC TGT GAT GAC CCC CGC TTC CAG GAC TCC	1205
Pro Lys Asp His Pro Leu Thr Cys Asp Asp Pro Arg Phe Gln Asp Ser	
580 585 590	
TCT TCC TCA AAG GCC CCT CCC CCC AGC CTT CCA AGC CCA TCC CGA CTC	1253
Ser Ser Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro Ser Arg Leu	
595 600 605 610	
CCG GGG CCC TCG GAC ACC CCG ATC CTC CCA CAA T AAGGATCCCT CGAG	1301
Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln	
615 620	

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 336 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Ser Arg Thr Ser Leu Leu Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp
1 5 10 15
Leu Gln Glu Gly Ser Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile
20 25 30
His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys Gly Thr
35 40 45
Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg
50 55 60

Glu	Cys	Glu	Ser	Gly	Ser	Phe	Thr	Ala	Ser	Glu	Asn	His	Leu	Arg	His	
65					70					75					80	
Cys	Leu	Ser	Cys	Ser	Lys	Cys	Arg	Lys	Glu	Met	Gly	Gln	Val	Glu	Ile	
				85					90					95		
Ser	Ser	Cys	Thr	Val	Asp	Arg	Asp	Thr	Val	Cys	Gly	Cys	Arg	Lys	Asn	
			100					105					110			
Gln	Tyr	Arg	His	Tyr	Trp	Ser	Glu	Asn	Leu	Phe	Gln	Cys	Phe	Asn	Cys	
		115					120					125				
Ser	Leu	Cys	Leu	Asn	Gly	Thr	Val	His	Leu	Ser	Cys	Gln	Glu	Lys	Gln	
	130					135					140					
Asn	Thr	Val	Cys	Thr	Cys	His	Ala	Gly	Phe	Phe	Leu	Arg	Glu	Asn	Glu	
145					150					155					160	
Cys	Val	Ser	Cys	Ser	Asn	Cys	Lys	Lys	Ser	Leu	Glu	Cys	Thr	Lys	Leu	
				165					170					175		
Cys	Leu	Pro	Gln	Ile	Glu	Asn	Val	Lys	Gly	Thr	Glu	Asp	Ser	Gly	Thr	
			180					185					190			
Thr	Ala	Gly	Ala	Gly	Pro	Arg	Cys	Arg	Pro	Ile	Asn	Ala	Thr	Leu	Ala	
		195					200					205				
Val	Glu	Lys	Glu	Gly	Cys	Pro	Val	Cys	Ile	Thr	Val	Asn	Thr	Thr	Ile	
	210					215					220					
Cys	Ala	Gly	Tyr	Cys	Pro	Thr	Met	Thr	Arg	Val	Leu	Gln	Gly	Val	Leu	
225					230					235					240	
Pro	Ala	Leu	Pro	Gln	Val	Val	Cys	Asn	Tyr	Arg	Asp	Val	Arg	Phe	Glu	
				245					250					255		
Ser	Ile	Arg	Leu	Pro	Gly	Cys	Pro	Arg	Gly	Val	Asn	Pro	Val	Val	Ser	
			260					265					270			
Tyr	Ala	Val	Ala	Leu	Ser	Cys	Gln	Cys	Ala	Leu	Cys	Arg	Arg	Ser	Thr	
		275					280					285				
Thr	Asp	Cys	Gly	Gly	Pro	Lys	Asp	His	Pro	Leu	Thr	Cys	Asp	Asp	Pro	
	290					295					300					
Arg	Phe	Gln	Asp	Ser	Ser	Ser	Ser	Lys	Ala	Pro	Pro	Pro	Ser	Leu	Pro	
305					310					315					320	
Ser	Pro	Ser	Arg	Leu	Pro	Gly	Pro	Ser	Asp	Thr	Pro	Ile	Leu	Pro	Gln	
				325					330				335			

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 6 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Ala Gly Ala Ala Pro Gly  
1 5

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 4 amino acids  
    (B) TYPE: amino acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Ala Gly Ala Gly  
1

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 30 base pairs  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TTTTCTCGAG ATGGCTACAG GTAAGCGCCC

30

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 39 base pairs  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

ACCTGGGGCA GCACCGGCAC AGGAGACACA CTCGTTTTTC

39

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 42 base pairs  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

TGTGCCGGTG CTGCCCCAGG TTGCCCAGAA TGCACGCTAC AG

42

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 36 base pairs  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TTTGTGGATCC TTAAGATTTG TGATAATAAC AAGTAC 36

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CCGTGGACCA GCACCAGCAC AGGAGACACA CTCGTTTTTC 39

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

TGTGCTGGTG CTGGTCCACG GTGCCGCCCC ATCAAT 36

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

TTTGTGGATCC TTATTGTGGG AGGATCGGGG TG 32

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

TTTTAGATCT CTTCTTGCAC AGTGGAC 27

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 21 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

TGTGGTGCCT GAGTCCTCAG T

21

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 41 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

ACTGAGGACT CAGGCACCAC AGCCGGTGCT GCCCCAGGTT G

41

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 29 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

TTTTTCTAGA GAAGCAGCAG CAGCCCATG

29

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 75 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

TTTTCCACAG CCAGGGTGGC ATTGATGGGG CGGCACCGTG GACCAGCACC AGCTGTGGTG

60

CCTGAGTCCT CAGTG

75

## CLAIMS

1. A hybrid protein comprising two coexpressed amino acid sequences forming a dimer, each comprising:

a) at least one amino acid sequence selected from the group consisting of a homomeric receptor, a chain of a heteromeric receptor, a ligand, and fragments thereof which retain the ligand-receptor binding capability; and

b) a subunit of a heterodimeric proteinaceous hormone, or fragments thereof which retain the ability of the subunit to form a heterodimer with other subunits thereof;

wherein sequences (a) and (b) are bonded directly or through a peptide linker, and in which the sequence (b) in each of said two coexpressed sequences are capable of aggregating to form a dimer complex.

2. A hybrid protein in accordance with claim 1, wherein said sequence (a) is selected from the group consisting of TBP1, TBP2 or fragments thereof still containing the ligand binding domain; the extracellular domain of the IFN $\alpha/\beta$  receptor or the IFN $\gamma$  receptor; a gonadotropin receptor or extracellular fragments thereof; antibody light chains or fragments thereof, optionally associated with the respective heavy chains; antibody heavy chains or fragments thereof; antibody Fab domains; and IL-6, IFN- $\beta$ , TPO or fragments thereof.

3. A hybrid protein in accordance with claim 1, wherein said sequence (b) is selected from the group consisting of subunits of hCG, FSH, LH, TSH or inhibin, and fragments thereof.

4. A hybrid protein in accordance with claim 1, wherein sequence (a) is linked to the amino terminus of sequence (b).



5. A hybrid protein in accordance with claim 1, wherein sequence (a) is linked to the carboxy terminus of sequence (b).

6. A hybrid protein in accordance with claim 1, wherein said two coexpressed amino acid sequences each include the sequence for TBP1 or the fragment thereof corresponding to amino acid residues 20-161 or 20-190 of TBP1, as sequence (a) and the respective  $\alpha$  and  $\beta$  subunits of hCG or fragments thereof, as sequence (b).

7. A hybrid protein in accordance with claim 1, wherein said two coexpressed amino acid sequences each include the extracellular domain of a gonadotropin receptor as sequence (a) and the respective  $\alpha$  and  $\beta$  subunits of a gonadotropin as sequence (b).

8. A hybrid protein in accordance with claim 7, wherein said sequence (a) is the FSH receptor extracellular domain and sequence (b) is a subunit of FSH.

9. A hybrid protein in accordance with claim 7, wherein said sequences (a) and (b) are linked with a peptide linker.

10. A hybrid protein in accordance with claim 9, wherein said peptide linker has an enzyme cleavage site.

11. A hybrid protein in accordance with claim 10, wherein said enzyme cleavage site is a thrombin cleavage site.

12. A hybrid protein in accordance with claim 10, wherein said enzyme cleavage site is recognized and cleaved by an enzyme which is found in the ovary.

13. A hybrid protein in accordance with claim 9, wherein said peptide linker serves as a flexible hinge.

14. A hybrid protein in accordance with claim 1, wherein one or more covalent bonds between the two subunits (b) are added.

15. A DNA molecule encoding a hybrid protein in accordance with claim 1.

16. An expression vector containing a DNA molecule in accordance with claim 15.

17. A host cell containing an expression vector in accordance with claim 16 and capable of expressing said hybrid protein.

18. A method for producing hybrid protein comprising culturing a host cell in accordance with claim 17 and recovering the hybrid protein expressed thereby.

19. A pharmaceutical composition comprising a hybrid protein in accordance with claim 1 and a pharmaceutically acceptable carrier and/or excipient.

20. A method for inducing follicular maturation, comprising administering a pharmaceutical composition comprising the hybrid protein of claim 8 to a subject in need thereof.

#### ABSTRACT OF THE DISCLOSURE

A hybrid protein includes two coexpressed amino acid sequences forming a dimer. Each sequence contains the binding portion of a receptor, such as TBP1 or TBP2, or a ligand, such as IL-6, IFN- $\beta$  and TPO, linked to a subunit of a heterodimeric proteinaceous hormone, such as hCG. Each coexpressed sequence contains a corresponding hormone subunit so as to form a heterodimer upon expression. Corresponding DNA molecules, expression vectors and host cells are also disclosed as are pharmaceutical compositions and a method of producing such proteins.

**Combined Declaration for Patent Application and Power of Attorney**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name; and that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

HYBRID PROTEINS the specification of which (check one)

[ ] is attached hereto;

[XX] was filed in the United States under 35 U.S.C. §111 on 20 February 1997, as USSN \_\_\_\_\_\*; or

[ ] was/will be filed in the U.S. under 35 U.S.C. §371 by entry into the U.S. national stage of an international (PCT) application, PCT/\_\_\_\_\_; filed \_\_\_\_\_, entry requested on \_\_\_\_\_\*; national stage application received USSN \_\_\_\_\_\*; §371/§102(e) date \_\_\_\_\_\* (\*if known),

and was amended on \_\_\_\_\_ (if applicable).

(include dates of amendments under PCT Art. 19 and 34 if PCT)

I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above; and I acknowledge the duty to disclose to the Patent and Trademark Office (PTO) all information known by me to be material to patentability as defined in 37 C.F.R. § 1.56.

I hereby claim foreign priority benefits under 35 U.S.C. §§ 119, 365 of any prior foreign application(s) for patent or inventor's certificate, or prior PCT application(s) designating a country other than the U.S., listed below with the "Yes" box checked and have also identified below any such application having a filing date before that of the application on which priority is claimed:

(Number)	(Country)	(Day Month Year Filed)	[ ] YES	[ ] NO
(Number)	(Country)	(Day Month Year Filed)	[ ] YES	[ ] NO
(Number)	(Country)	(Day Month Year Filed)	[ ] YES	[ ] NO

I hereby claim the benefit under 35 U.S.C. § 120 of any prior U.S. non-provisional Application(s) or prior PCT Application(s) designating the U.S. listed below, or under § 119(e) of any prior U.S. provisional applications listed below, and, insofar as the subject matter of each of the claims of this application is not disclosed in such U.S. or PCT application in the manner provided by the first paragraph of 35 U.S.C. §112, I acknowledge the duty to disclose to the PTO all information as defined in 37 C.F.R. §1.56(a) which occurred between the filing date of the prior application and the national filing date of this application:

<u>60/011,936</u>	<u>20 February 1996</u>	<u>Abandoned</u>
(Application Serial No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)
(Application Serial No.)	(Day Month Year Filed)	(Status: patented, pending, abandoned)

I hereby appoint the following attorneys, with full power of substitution, association, and revocation, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith.

SHERIDAN NEIMARK, REG. NO. 20,520 - ROGER L. BROWDY, REG. NO. 25,618 - ANNE M. KORNBAU, REG. NO. 25,884  
 NORMAN J. LATKER, REG. NO. 19,963 - IVER P. COOPER, REG. NO. 28,005 - ALLEN C. YUN, REG. NO. 37,971\*  
 NICK S. BROMER, REG. NO. 33,478 - \* Patent Agent

ADDRESS ALL CORRESPONDENCE TO

**BROWDY AND NEIMARK, P.L.L.C.**  
419 Seventh Street, N.W.  
Washington, D.C. 20004

DIRECT ALL TELEPHONE CALLS TO:

**BROWDY AND NEIMARK**  
 (202) 628-5197




The undersigned hereby authorizes the U.S. Attorneys or Agents named herein to accept and follow instructions from INSTITUTO FARMACOLOGICA SERONO S.P.A. as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. Attorney or Agent and the undersigned. In the event of a change of the persons from whom instructions may be taken, the U.S. Attorneys or Agents named herein will be so notified by the undersigned.


Title: HYBRID PROTEINS

U.S. Application filed 20 February 1997, Serial No. \_\_\_\_\_

PCT Application filed \_\_\_\_\_, Serial No. \_\_\_\_\_

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. §1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

FULL NAME OF FIRST INVENTOR <u>Robert K. CAMPBELL</u> 1-00	INVENTOR'S SIGNATURE 	DATE 4 June 1997
RESIDENCE <u>Wrentham, Massachusetts, U.S.A. MA</u>		CITIZENSHIP American
POST OFFICE ADDRESS 25 Meadowbrook Drive, Wrentham, Massachusetts, U.S.A.		
FULL NAME OF SECOND JOINT INVENTOR <u>Bradford A. JAMESON</u> 2-00	INVENTOR'S SIGNATURE 	DATE 4 June 97
RESIDENCE <u>Milton, Massachusetts, U.S.A. MA</u>		CITIZENSHIP American
POST OFFICE ADDRESS 76 Robbins Street, Milton, Massachusetts, U.S.A.		
FULL NAME OF THIRD JOINT INVENTOR <u>Scott C. CHAPPEL</u> 3-00	INVENTOR'S SIGNATURE 	DATE June 4, 1997
RESIDENCE <u>Milton, Massachusetts, U.S.A. MA</u>		CITIZENSHIP American
POST OFFICE ADDRESS 125 Canton Avenue, Milton, Massachusetts, U.S.A.		
FULL NAME OF FOURTH JOINT INVENTOR	INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP
POST OFFICE ADDRESS		
FULL NAME OF FIFTH JOINT INVENTOR	INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP
POST OFFICE ADDRESS		
FULL NAME OF SIXTH JOINT INVENTOR	INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP
POST OFFICE ADDRESS		
FULL NAME OF SEVENTH JOINT INVENTOR	INVENTOR'S SIGNATURE	DATE
RESIDENCE		CITIZENSHIP
POST OFFICE ADDRESS		



XhoI hGH Signal Sequence hGH Intron

TCGAG ATG GCT ACA G GAAAGCGCCCTAAATCCCTTTGGGCACATGTCTCTGAGGGAGAGGCAGCGACCTGTAGATGGGACGGGGCACTAACCCCTCAGGTTTGGGGCTTCT

► Met Ala Thr

GAATGTGAGTATCGCCATGTAAGCCCAATTTGGCCCAATCTCAGAAAGCTCCTGGTCCCTGGAGGGATGGAGAGAGAAAAACAACAGCTCCTGGAGCAGGGAGAGTGCCTGGCCTCTTGCTCTC

CGGCTCCCTCTGTTGCCCTCTGGTTTCTCCCCAGGC TCC CGG ACG TCC CTG CTC CTG GCT TTT GGC CTG CTC TGC CTG CCC TGG CTT

► Ser Arg Thr Ser Leu Leu Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp Leu

+20 Asp of Processed TBP1

CAA GAG GGC AGT GGC GAT AGT GTG TGT CCC CAA GGA AAA TAT ATC CAC CCT CAA AAT AAT TCG ATT TGC TGT ACC AAG TGC CAC AAA GGA

► Gln Glu Gly Ser Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys Gly

ACC TAC TTG TAC AAT GAC TGT CCA GGC CCG GGG CAG GAT ACG GAC TGC AGG GAG TGT GAG AGC GGC TCC TTC ACC GCT TCA GAA AAC CAC CTC

► Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu

AGA CAC TGC CTC AGC TGC TCC AAA TGC CGA AAG GAA ATG GGT CAG GTG GAG ATC TCT TCT TGC ACA GTG GAC CGG GAC ACC GTG TGT GGC TGC

► Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys

AGG AAG AAC CAG TAC CGG CAT TAT TGG AGT GAA AAC CTT TTC CAG TGC TTC AAT TGC AGC CTC TGC CTC AAT GGG ACC GTG CAC CTC TCC TGC

► Arg Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys

Linker

CAG GAG AAA CAG AAC ACC GTG TGC ACC TGC CAT GCA GGT TTC TTT CTA AGA GAA AAC GAG TGT GTC TCC TGT GGC GGT GCT GCC CCA GGT

► Gln Glu Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ala Gly Ala Ala Pro Gly

+7 Cys of hCG alpha

TGC CCA GAA TGC ACG CTA CAG GAA AAC CCA TTC TTC TCC CAG CCG GGT GCC CCA ATA CTT CAG TGC ATG GGC TGC TGC TTC TCT AGA GCA TAT

► Cys Pro Glu Cys Thr Leu Gln Glu Asn Pro Phe Phe Ser Gln Pro Gly Ala Pro Ile Leu Gln Cys Met Gly Cys Cys Phe Ser Arg Ala Tyr

CCC ACT CCA CTA AGG TCC AAG AAG ACG ATG TTG GTC CAA AAG AAC GTC ACC TCA GAG TCC ACT TGC TGT GTA GCT AAA TCA TAT AAC AGG GTC

► Pro Thr Pro Leu Arg Ser Lys Lys Thr Met Leu Val Gln Lys Asn Val Thr Ser Glu Ser Thr Cys Cys Val Ala Lys Ser Tyr Asn Arg Val


ACA GTA ATG GGG GGT TTC AAA GTG GAG AAC CAC ACG GCG TGC CAC TGC AGT ACT TGT TAT TAT CAC AAA TCT TAA G

► Thr Val Met Gly Gly Phe Lys Val Glu Asn His Thr Ala Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser ...

Bam HI

Figure 1 (a)

TBP(20-161)-hCG $\alpha$  FUSION CONSTRUCT



**hGH Signal Sequence** **hGH Intron**

**XhoI**

CTCGAG ATG GCT ACA G GTAAGCGCCCTAAATCCCTTTGGGCACATGTGTCTGAGGGAGAGGGCAGCGACCTGTAGATGGGACGGGGGCACTACCCCTCAGGTTTGGG

► Met Ala Thr

GCTTCIGATGTGAGTATCGCCATGTAAGCCCATTTGGCCATCTCAGAAAGCTCCTGGTCCCTGGAGGGATGGAGAGAGTAAACAAACAGCTCCTGGAGCAGGGAGAGTGGTGGC

CTCTTGCCTCTCCGGCTCCCTCTGTGCTCCCTCTGTTTCTCCCCAGGC TCC CGG ACG TCC CTG CTC CTG GCT TTT GGC CTG CTC TGC CTG

► Ser Arg Thr Ser Leu Leu Leu Ala Phe Gly Leu Leu Cys Leu

+20 Asp of Processed TBP1

CCC TGG CTT CAA GAG GGC AGT GCC GAT AGT GTG TGT CCC CAA GGA AAA TAT ATC CAC COT CAA AAT AAT TCG ATT TGC TGT ACC

► Pro Trp Leu Gln Glu Gly Ser Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr

AAG TGC CAC AAA GGA ACC TAC TTG TAC AAT GAC TGT CCA GGC CCG GGG CAG GAT ACG GAC TGC AGG GAG TGT GAG AGC GGC TCC TTC ACC

► Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr

GCT TCA GAA AAC CAC CTC AGA CAC TGC CTC AGC TGC TCC AAA TGC CGA AAG GAA ATG GGT CAG GTG GAG ATC TCT TCT TGC ACA GTG GAC

► Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp

CGG GAC ACC GTG TGT GGC TGC AGG AAG AAC CAG TAC CGG CAT TAT TGG AGT GAA AAC CTT TTC CAG TGC TTC AAT TGC AGC CTC TGC CTC

► Arg Asp Thr Val Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu

AAT GGG ACC GTG CAC CTC TCC TGC CAG GAG AAA CAG AAC ACC GTG TGC ACC TGC CAT GCA GGT TTC TTT CTA AGA GAA AAC GAG TGT GTC

► Asn Gly Thr Val His Leu Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val

Linker +7 Pro of hCG beta

TCC TGT GCT GGT GCT GGT CCA CGG TGC CGC CCC ATC AAT GCC ACC CTG GCT GTG GAG AAG GAG GGC TGC CCC GTG TGC ATC ACC GTC

► Ser Cys Ala Gly Ala Gly Pro Arg Cys Arg Pro Ile Asn Ala Thr Leu Ala Val Glu Lys Glu Gly Cys Pro Val Cys Ile Thr Val

AAC ACC ACC ATC TGT GCC GGC TAC TGC CCC ACC ATG ACC CGC GTG CTG CAG GGG GTC CTG CCG GCC CTG CCT CAG GTG GTG TGC AAC TAC

► Asn Thr Thr Ile Cys Ala Gly Tyr Cys Pro Thr Met Thr Arg Val Leu Gln Gly Val Leu Pro Ala Leu Pro Gln Val Val Cys Asn Tyr

CGC GAT GTG CGC TTC GAG TCC ATC CGG CTC CCT GGC TGC CCG CGC GGC GTG AAC CCC GTG GTC TCC TAC GCC GTG GCT CTC AGC TGT CAA

► Arg Asp Val Arg Phe Glu Ser Ile Arg Leu Pro Gly Cys Pro Arg Gly Val Asn Pro Val Val Ser Tyr Ala Val Ala Leu Ser Cys Gln

TGT GCA CTC TGC CGC CGC AGC ACC ACT GAC TGC GGG GGT CCC AAG GAC CAC CCC TTG ACC TGT GAT GAC CCC CGC TTC CAG GAC TCC TCT


► Cys Ala Leu Cys Arg Arg Ser Thr Thr Asp Cys Gly Gly Pro Lys Asp His Pro Leu Thr Cys Asp Asp Pro Arg Phe Gln Asp Ser Ser

TCC TCA AAG GCC CCT CCC CCC AGC CTT CCA AGC CCA TCC CGA CTC CCG GGG CCC TCG GAC ACC CCG ATC CTC CCA CAA TTA

► Ser Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro Ser Arg Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln ... Bam HI

Figure 1 (b)

## TBP(20-161)-hCGβ FUSION CONSTRUCT



**XhoI**                      **hGH Signal Sequence**                      **hGH Intron**

TCGAG ATG GCT ACA G CTAAGCGCCCCCTAAATCCCTTTGGGCACAAATGTGTCCTGAGGGGAGAGGCAGCGACCTGTAGATGGGACGGGGGCACCTAACCCCTCAGGTTTGGGGCTTCT

► Met Ala Thr

GAATGTGAGTATCGCCATGTAAGCCAGTATTGGCCCAATCTCAGAAAGCTCCTGGTCCCTCGAGGGATGGAGAGAGAAAAACAAACAGCTCCTGGAGCAGGGAGAGTGCTGGCCTCTTGCTCTC

CGGCTCCCTCTGTTGGCCCTCGGTTTCTCCCGAGC TCC CGG ACG TCC CTG CTC CTG GCT TTT GGC CTG CTC TGC CTG CCC TGG CTT

► Ser Arg Thr Ser Leu Leu Leu Ala Phe Gly Leu Leu Cys Leu Pro Trp Leu

+20 Asp of processed TBP1

CAA GAG GGC AGT GCC GAT AGT GTG TGT CCC CAA GGA AAA TAT ATC CAC CCT CAA AAT AAT TCG ATT TGC TGT ACC AAG TGC CAC AAA GGA

► Gln Glu Gly Ser Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr Lys Cys His Lys Gly

ACC TAC TTG TAC AAT GAC TGT CCA GGC CCG GGG CAG GAT ACG GAC TGC AGG GAG TGT GAG AGC GGC TCC TTC ACC GCT TCA GAA AAC CAC CTC

► Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr Ala Ser Glu Asn His Leu

AGA CAC TGC CTC AGC TGC TCC AAA TGC CGA AAG GAA ATG GGT CAG GTG GAG ATC TCT TCT TGC ACA GTG GAC CGG GAC ACC GTG TGT GGC TGC

► Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp Arg Asp Thr Val Cys Gly Cys

AGG AAG AAC CAG TAC CGG CAT TAT TGG AGT GAA AAC CTT TTC CAG TGC TTC AAT TGC AGC CTC TGC CTC AAT GGG ACC GTG CAC CTC TCC TGC

► Arg Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu Asn Gly Thr Val His Leu Ser Cys

CAG GAG AAA CAG AAC ACC GTG TGC ACC TGC CAT GCA GGT TTC TTT CTA AGA GAA AAC GAG TGT GTC TCC TGT AGT AAC TGT AAG AAA AGC CTG

► Gln Glu Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val Ser Cys Ser Asn Cys Lys Lys Ser Leu

Linker +7 Cys of hCG alpha

GAG TGC ACG AAG TTG TGC CTA CCC CAG ATT GAG AAT GTT AAG GGC ACT GAG GAC TCA GGC ACC ACA GCC GGT GCT GCC CCA GGT TGC CCA

► Glu Cys Thr Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Ser Gly Thr Thr Ala Gly Ala Ala Pro Gly Cys Pro

GAA TGC ACG CTA CAG GAA AAC CCA TTC TTC TCC CAG CCG GGT GCC CCA ATA CTT CAG TGC ATG GGC TGC TGC TTC TCT AGA GCA TAT CCC ACT

► Glu Cys Thr Leu Gln Glu Asn Pro Phe Phe Ser Gln Pro Gly Ala Pro Ile Leu Gln Cys Met Gly Cys Cys Phe Ser Arg Ala Tyr Pro Thr

CCA CTA AGG TCC AAG AAG ACG ATG TTG GTC CAA AAG AAC GTC ACC TCA GAG TCC ACT TGC TGT GTA GCT AAA TCA TAT AAC AGG GTC ACA GTA

► Pro Leu Arg Ser Lys Lys Thr Met Leu Val Gln Lys Asn Val Thr Ser Glu Ser Thr Cys Cys Val Ala Lys Ser Tyr Asn Arg Val Thr Val


ATG GGG GGT TTC AAA GTG GAG AAC CAC ACG GCG TGC CAC TGC AGT ACT TGT TAT TAT CAC AAA TCT TAA GGATCCCTCGAG

► Met Gly Gly Phe Lys Val Glu Asn His Thr Ala Cys His Cys Ser Thr Cys Tyr Tyr His Lys Ser ... BamHI XhoI

Figure 2(a)  
TBP(20-190)-hCG $\alpha$  FUSION CONSTRUCT



08/204166



XhoI hGH Signal Sequence hGH Intron

CTCGAG ATG GCT ACA G GTAAAGCGCCCTAAATCCCTTTGGGCACAATGTGTCTGAGGGGAGAGGCGGACCTGTAGATGGGACGGGGCCACTAACCCCTCAGGTTTGGG

►Met Ala Thr

GCTTCTGATGTGAGTATCGCCATGTAAGCCAGTATTGGCCCAATCTCAGAAAGCTCCTGGTCCCTGGAGGGATGGAGAGAGAAAAACAAACAGCTCCTCGAGCAGGGAGAGTGTGGC

CTCTTGCTCTCCGGCTCCCTCTGTGTGCCCTCTGGTTTCCTCCCAAGGC TCC CGG ACG TCC CTG CTC CTG GCT TTT GGC CTG CTC TGC CTG

►Ser Arg Thr Ser Leu Leu Leu Ala Phe Gly Leu Leu Cys Leu

+20 Asp of Processed TBP1

CCC TGG CTT CAA GAG GGC AGT GCC GAT AGT GTG TGT CCC CAA GGA AAA TAT ATC CAC CCT CAA AAT AAT TCG ATT TGC TGT ACC

►Pro Trp Leu Gln Glu Gly Ser Ala Asp Ser Val Cys Pro Gln Gly Lys Tyr Ile His Pro Gln Asn Asn Ser Ile Cys Cys Thr

AAG TGC CAC AAA GGA ACC TAC TTG TAC AAT GAC TGT CCA GGC CCG GGG CAG GAT ACG GAC TGC AGG GAG TGT GAG AGC GGC TCC TTC ACC

►Lys Cys His Lys Gly Thr Tyr Leu Tyr Asn Asp Cys Pro Gly Pro Gly Gln Asp Thr Asp Cys Arg Glu Cys Glu Ser Gly Ser Phe Thr

GCT TCA GAA AAC CAC CTC AGA CAC TGC CTC AGC TGC TCC AAA TGC CGA AAG GAA ATG GGT CAG GTG GAG ATC TCT TCT TGC ACA GTG GAC

►Ala Ser Glu Asn His Leu Arg His Cys Leu Ser Cys Ser Lys Cys Arg Lys Glu Met Gly Gln Val Glu Ile Ser Ser Cys Thr Val Asp

CGG GAC ACC GTG TGT GGC TGC AGG AAG AAC CAG TAC CGG CAT TAT TGG AGT GAA AAC CTT TTC CAG TGC TTC AAT TGC AGC CTC TGC CTC

►Arg Asp Thr Val Cys Gly Cys Arg Lys Asn Gln Tyr Arg His Tyr Trp Ser Glu Asn Leu Phe Gln Cys Phe Asn Cys Ser Leu Cys Leu

AAT GGG ACC GTG CAC CTC TCC TGC CAG GAG AAA CAG AAC ACC GTG TGC ACC TGC CAT GCA GGT TTC TTT CTA AGA GAA AAC GAG TGT GTC

►Asn Gly Thr Val His Leu Ser Cys Gln Glu Lys Gln Asn Thr Val Cys Thr Cys His Ala Gly Phe Phe Leu Arg Glu Asn Glu Cys Val

TCC TGT AGT AAC TGT AAG AAA AGC CTG GAG TGC ACG AAG TTG TGC CTA CCC CAG ATT GAG AAT GTT AAG GGC ACT GAG GAC TCA GGC ACC

►Ser Cys Ser Asn Cys Lys Lys Ser Leu Glu Cys Thr Lys Leu Cys Leu Pro Gln Ile Glu Asn Val Lys Gly Thr Glu Asp Ser Gly Thr

Linker +7 Pro of beta

ACA GCT GGT GCT GGT CCA CGG TGC CGC CCC ATC AAT GCC ACC CTG GCT GTG GAG AAG GAG GGC TGC CCC GTG TGC ATC ACC GTC AAC

►Thr Ala Gly Ala Gly Pro Arg Cys Arg Pro Ile Asn Ala Thr Leu Ala Val Glu Lys Glu Gly Cys Pro Val Cys Ile Thr Val Asn

ACC ACC ATC TGT GCC GGC TAC TGC CCC ACC ATG ACC CGC GTG CTG CAG GGG GTC CTG CCG GCC CTG CCT CAG GTG GTG TGC AAC TAC CGC

►Thr Thr Ile Cys Ala Gly Tyr Cys Pro Thr Met Thr Arg Val Leu Gln Gly Val Leu Pro Ala Leu Pro Gln Val Val Cys Asn Tyr Arg

GAT GTG CGC TTC GAG TCC ATC CGG CTC CCT GGC TGC CCG CGC GGC GTG AAC CCC GTG GTC TCC TAC GCC GTG GCT CTC AGC TGT CAA TGT

►Asp Val Arg Phe Glu Ser Ile Arg Leu Pro Gly Cys Pro Arg Gly Val Asn Pro Val Val Ser Tyr Ala Val Ala Leu Ser Cys Gln Cys

GCA CTC TGC CGC CGC AGC ACC ACT GAC TGC GGG GGT CCC AAG GAC CAC CCC TTG ACC TGT GAT GAC CCC CGC TTC CAG GAC TCC TCT TCC


►Ala Leu Cys Arg Arg Ser Thr Thr Asp Cys Gly Gly Pro Lys Asp His Pro Leu Thr Cys Asp Asp Pro Arg Phe Gln Asp Ser Ser Ser

TCA AAG GCC CCT CCC CCC AGC CTT CCA AGC CCA TCC CGA CTC CCG GGG CCC TCG GAC ACC CCG ATC CTC CCA CAA TAA GGATCCCTCGAG


►Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro Ser Arg Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln ... BamHI XhoI

Figure 2 (b)  
TBP(20-190)-hCGβ FUSION CONSTRUCT


Figure 4. CHO cell expressed TBP-hCG(20-190) inhibits TNF $\alpha$ -induced cytotoxicity on BT-20 cells



084804166



**Figure 5.** COS cell expressed TBP-hCG(20-190) inhibits TNF $\alpha$ -induced cytotoxicity on BT-20 cells



**Figure 6.** Affinity purified CHO cell expressed TBP-hCG(20-161) inhibits TNF $\alpha$ -induced cytotoxicity on BT-20 cells

